



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Title: The Need for Eccentric Speed: A Narrative Review of the Effects of Accelerated Eccentric Actions During Resistance-Based Training

Author(s): Handford, Matthew, Bright, Thomas, Munday, Peter, Lake, Jason, Theis, Nicola, Hughes Jonathan

Copyright, publisher and additional information: © The Author(s), under exclusive licence to Springer Nature Switzerland AG 2022

DOI: 10.1007/s40279-022-01686-z

Reference: Handford, M.J., Bright, T.E., Mundy, P. et al. The Need for Eccentric Speed: A Narrative Review of the Effects of Accelerated Eccentric Actions During Resistance-Based Training. *Sports Med* (2022). <https://doi.org/10.1007/s40279-022-01686-z>

Accepted 30 March 2022

Published 10 May 2022

DOI <https://doi.org/10.1007/s40279-022-01686-z>

24 **Title:** The need for eccentric speed: A narrative review of the effects of accelerated eccentric actions during
25 resistance-based training.

26 **Running Header:** A narrative review of accelerated eccentrics.

27

28

29

30 **Authors**

31 Matthew J. Handford ¹, Thomas E. Bright ^{1,4}, Peter Mundy ², Jason Lake ^{3,5}, Nicola Theis ¹, Jonathan D. Hughes
32 ¹

33

34

35 ¹ School of Sport and Exercise, University of Gloucestershire, Gloucester, United Kingdom

36 ² Faculty of Health and Life Sciences, Coventry University, Coventry, United Kingdom

37 ³ Chichester Institute of Sport, Nursing, and Allied Health, University of Chichester, Chichester, United
38 Kingdom

39 ⁴ School of Sport, Health and Wellbeing, Plymouth Marjon University, Plymouth, United Kingdom

40 ⁵ School of Medical and Health Sciences, Edith Cowan University, Joondalup, Australia

41

42

43

44 Corresponding author: Matthew J Handford. Email: s1507429@glos.ac.uk

45

46 Ethics Declarations: Not applicable.

47 Funding: No funding was received to complete the current research.

48 Conflicts of interest: The author declares no conflict of interest.

49 Ethics approval: Ethical approval for publication of this review was not required. However, the authors declare
50 that the findings of this review are presented honestly and without any fabrication, falsification, or inappropriate
51 data manipulation.

52 Consent to participate: Not applicable.

53 Consent for publication: All authors declare consent for publication.

54 Availability of data and materials: Data sharing is not applicable to this article as no datasets were generated or
55 analysed during the current review.

56 Code availability: Not applicable

57 Author contribution: All authors declare that they contributed to the completion of this paper.

58 Declaration: All authors read and approved the final version

59 **Abstract**

60 Eccentric training, as a method to enhance athletic performance, is a topic of increasing interest to both
61 practitioners and researchers. However, there is limited data regarding the effects of performing eccentric actions
62 of an exercise at increased velocities. This narrative review aimed to provide greater clarity for eccentric methods
63 and classification with regard to temporal phases of exercises. To achieve the object of the review, key terms were
64 searched using PubMed, SPORTDiscus and Google Scholar databases between March and April 2021 within the
65 years of 1950-2021. Search terms included: ('fast eccentric'), ('fast velocity eccentric'), ('dynamic eccentric'),
66 ('accentuated eccentric loading'), ('isokinetic eccentric'), analysing both the acute and chronic effects of
67 accelerated eccentric training on human participants. Of the 26 studies which met inclusion criteria, it was
68 identified that completing eccentric tempos of <2s can increase subsequent concentric one repetition maximum
69 performance, velocity, and power, compared to >4s tempos. Durations of >4s tempo increase time under tension
70 (TUT); whilst reduced tempos allow for greater volume to be completed. Greater TUT leads to larger
71 accumulation of blood lactate, growth hormone and testosterone, when volume is matched to that of the reduced
72 tempos. Overall, evidence supports <2s duration eccentric actions to improve subsequent concentric performance.
73 There is no clear difference between using eccentric tempos of 2–6s if the aim is to increase hypertrophic response
74 and strength. Future research should analyse performing eccentric actions at greater velocities or reduced time
75 durations to determine more factors such as strength response. Tempo studies should aim to complete the same
76 TUT for protocols to determine measures for hypertrophic response.

77

78

79 **Key Points**

80 The term of Accelerated Eccentrics should be reintroduced to define eccentric actions that are performed at
81 increased velocities with the addition of specific verbal cues.

82 There is no clear difference between <2s versus >4s eccentric tempos and <60°s⁻¹ versus >180°s⁻¹ eccentric
83 isokinetic dynamometry for increasing muscle mass in a longitudinal training programme.

84 There is a need for more academic research to view the effects of accelerated eccentrics to determine how it
85 impacts both the acute and longitudinal response to training.

86 1.0 Introduction

87 Eccentric training as a method to enhance athletic performance is a topic constantly under review as researchers
88 aim to understand the physical capabilities that these muscle actions can elicit. There is a gap in the knowledge
89 base when comparing the eccentric to the concentric side of the force-velocity curve [1,2]. It is widely accepted
90 that during multi-and single-joint actions, eccentric muscle lengthening forces can reach 100-150% of a concentric
91 one repetition maximum (1RM) or 20-60% greater than concentric actions [3–8]. The emergence of this
92 understanding has led academics and coaches to create, test, and implement training methods to incorporate this
93 mode of training. Primarily, methods have concentrated on increasing the mass utilised to increase eccentric forces
94 illustrated via supramaximal loads through methods such as accentuated eccentric loading (AEL) [9–11].

95 The training procedure of accelerated eccentrics is often overlooked despite regular discussion in the literature
96 (Table 1) [5,12–17], this centres on increasing the acceleration (eccentric velocity) of any movement, though this
97 is not fully investigated [18]. Eccentric force-velocity relationships differ to concentric muscle actions [18,19]
98 first devised by Hill’s muscle model [20]. During eccentric muscle lengthening, higher forces are attained
99 alongside high velocity [13,14,16,20–28]. Though, this statement only holds true to a certain degree, for example,
100 if tension (force and velocity) is too great then an individual would not be able to produce enough force to
101 decelerate and would continue to displace through a motion [23,27,29]. The enhancement in force production is
102 thought to be achieved by an increase in the proportion of cross bridges or an increase in the average force per
103 cross bridge or both [22]. The giant protein Titin has also been theorised to play a role by storing elastic energy
104 in the I-band region [3,12,22]. Evidence demonstrates that the key role for Titin in the lasting force enhancement
105 during lengthening actions is down to increases in Titin stiffness termed “passive force enhancement” [30–33].
106 Therefore, it is now more prudent to understand that it is not only cross-bridges, but also Titin, that contribute to
107 enhanced force levels during lengthening actions as Titin demonstrably enhances stiffness during all other phases
108 of an eccentric contraction.

109 Biomechanically, utilising greater eccentric velocities may provide training performance enhancements. If the
110 aim is to increase the acceleration of a known mass over a set range of a motion than a greater force is required to
111 be able to accelerate a mass at a greater rate ($F = m \times a$). When observing kinetic energy ($ke = m \times v^2$), increasing
112 the velocity has a greater effect on ke than enhancing mass [13,16,37]. This can be seen when employing drop
113 and depth jumps (shock method) as the height of the platform is increased potential energy is augmented as there
114 is a larger distance to accelerate through. In doing so this potential energy is transferred into ke once a person

115 steps off a platform and begins to free fall [13,38,39]. This asserts that to increase the acceleration of a movement
116 force must be enhanced or the duration at which force is applied can increase, therefore a coach can increase the
117 aforementioned metrics without the need to increase the mass ensuring duration remains the same.

118 Momentum is defined as mass in motion and again can be enhanced by increasing velocity or mass. Therefore, if
119 an object is travelling at a greater velocity then an increased force would be required to decelerate it [40]. This
120 can also be altered by an individual performing voluntary actions such as pulling the barbell down to increase
121 eccentric acceleration. Subsequently, some authors have investigated increasing eccentric velocity during
122 resistance-based training, this method though is performed as an isoinertial action and will achieve a greater
123 eccentric tension as the velocity of the action is increased. This method would apply a far simpler process than
124 weight releasers [41–43] or specialized equipment [44] found during supramaximal loaded AEL studies.
125 Subsequently, being more pragmatic for coaches to apply greater emphasis on increased eccentric velocity
126 exercises, though there is little evidence to support this claim.

127 Eccentric training has been recognized as a valid training method for performance enhancement [14,43]; however,
128 there is a distinct lack of consistency in the terms used to describe the duration and velocity of eccentric training
129 in the research. Furthermore, there is limited research into the area of accelerated eccentric training, thus a review
130 of the current topic is warranted. The aims of this narrative review is to firstly, examine the available literature
131 pertaining to accelerated eccentric training and the application of this modality to provide a greater understanding
132 of the adaptations that are evidenced to occur from accelerated eccentrics and secondly, to provide more consistent
133 and accurate terminology for practitioners to use.

134

135

136

137 *****INSERT TABLE 1 HERE*****

138

139

140

141

142 **2.0 Methods**

143 **2.1 Database Search**

144 A literature search was conducted from 02/03/2021 with the last search being performed on 21/04/2021 with the
145 years of research obtained from 1950-2021. Pubmed and SPORTDiscus search engines were utilised whereby
146 title, abstract and key words were viewed. Google Scholar was searched using the advanced search section.
147 Furthermore, reference lists from the identified studies via the database searchers were also examined to discover
148 any additional research that could be used in the current review. The following search terms were included: ('fast
149 eccentric'), ('eccentric velocity'), ('dynamic eccentric'), ('overspeed eccentric'), ('accentuated eccentric
150 loading'), ('isokinetic eccentric'), ('lengthening velocity'), ('deceleration training'), ('fast stretch shortening
151 cycle'). The study analysed both the acute and chronic effects of accelerated eccentric training on human
152 participants. Research studies that only utilised one eccentric aforementioned value (duration or velocity) would
153 be utilised to justify results but not used in the analysis (Table 2).

154 **2.2 Inclusion and Exclusion Criteria**

155 Initially, after studies were collated if they did not complete the same Concentric methods they were excluded as
156 this would not allow comparisons to be made across protocols. In addition to this the studies must further have
157 completed the inclusion criteria seen in Table 2.

158

159

160

**** INSERT TABLE 2 HERE ****

161

162

163

164

165 **3.0 Results**

166 **3.1 Literature Search Results**

167 Of the 987 studies found during the database search, 49 met the inclusion criteria, these were then further analysed
168 via a modified Downs-Black assessment [88] (Figure 1). Two separate modified Downs-Black assessments were
169 undertaken as the current review analysed both the acute and chronic effects of accelerated eccentric training.
170 Two researchers conducted the modified Downs-Black assessments, if studies achieved a score of 10 and below
171 during the modified Downs-Black assessment, they were removed from the analysis to better improve the current
172 studies bias. After the modified Downs-Black analysis, 26 studies were selected for further review. Acute results
173 of tempo studies and isokinetic dynamometer can be found in Table 3 and 4, respectively. The chronic effects of
174 tempo and isokinetic dynamometry can be found in Table 5 and 6, respectively.

175

176

177

***** INSERT FIGURE 1 HERE *****

178

179

180 **3.2: Tables of Findings**

181

182

**** INSERT TABLE 3 HERE ****

183

184

**** INSERT TABLE 4 HERE ****

185

186

187

**** INSERT TABLE 5 HERE ****

188

189

190

**** INSERT TABLE 6 HERE ****

191

192 **4.0 Discussion**

193 **4.1 Eccentric Torque Velocity Relationship**

194 The results of this review indicate that during eccentric IKD greater forces can be achieved when angular velocities
195 $>180^{\circ}\cdot s^{-1}$ is performed when compared to $<180^{\circ}\cdot s^{-1}$ [14,16,78,99,100] and some studies indicated that there are
196 no difference in torque value when values of $180^{\circ}\cdot s^{-1}$ and above are performed [25,54,95,101] (Table 4). The
197 chronic effects of utilising eccentric IKD identified that 180 and $210^{\circ}\cdot s^{-1}$ angular velocity value groups improved
198 all tested angular velocity metrics [53,55,56,58], whereas the 20 and $30^{\circ}\cdot s^{-1}$ groups did not [53,55,56,58] (Table
199 6). These findings are in agreement with Kellis and Baltzopoulos' [52] review which explained that the eccentric
200 IKD is not velocity specific. The increase in torque value results are expected as eccentric torque values are
201 enhanced or maintained when there is an increase in velocity [23,102–104]. Therefore, participants will increase
202 force production and overall strength capacity of the muscle through physiological factors e.g., increasing motor
203 unit recruitment [3,102]. When eccentric actions are performed maximally a greater muscular action will occur,
204 therefore, an increase in motor unit recruitment will arise [102]. This with the addition of Titin being thought to
205 enhance force production a greater muscle action could achieve subsequent enhanced physiological adaptations
206 occur [30–33]. This concurs with Verkoshansky and Siff [13] who state faster actions have a better transfer to
207 slow actions than vice versa.

208 Farthing and Chilibeck [56] found chronic $180^{\circ}\cdot s^{-1}$ eccentric IKD increased concentric peak torque at 180 and
209 $30^{\circ}\cdot s^{-1}$. Paddon-Jones [58] observed $180^{\circ}\cdot s^{-1}$ eccentric IKD increased concentric peak torque at $180^{\circ}\cdot s^{-1}$ but neither
210 $180^{\circ}\cdot s^{-1}$ or $30^{\circ}\cdot s^{-1}$ increased concentric peak torque testing values (Table 6). Previous findings are mixed for mode
211 specific IKD and very little research has viewed two different eccentric torque values in relation to increasing
212 concentric strength (Table 6). From these results it can be hypothesised $180^{\circ}\cdot s^{-1}$ eccentric IKD could enhance
213 concentric IKD torque values greater than $<180^{\circ}\cdot s^{-1}$ IKD scores however, future research is required to define a
214 definitive conclusion.

215 **4.2 Eccentric Tempo Acute Response**

216 Acute studies investigating the impact of reduced durations of movement on concentric power and/or velocity
217 found a <2 s eccentric duration significantly increased power and velocity [66,81,89–91] in subsequent concentric
218 performance (Table 3). Schoenfeld [105] commented that slower tempos lead to less concentric force, power and
219 reduced stretch shortening cycle (SSC) via a reduced recruitment of type IIx muscle fibres. Carzoli et al. [89]

220 highlighted through completion of faster eccentric actions during both back squat and bench press with 60 and
221 80% of 1RM a subsequent enhancement of concentric average and peak velocities (Table 3). It can be suggested
222 that these findings occur from a better response of SSC [106] and faster eccentric muscle actions achieve a greater
223 effect of type IIa and IIx muscle fibres [78] and motor unit recruitment. However, only one study in the current
224 review analysed eccentric kinematics results [81]. Therefore, future research must address this method so a more
225 definitive conclusion can be addressed.

226 Several acute studies investigating the impact of tempo on strength values have indicated that faster eccentric
227 tempos elicit greater strength values throughout a whole exercise movement [72,93,97] (Table 3). In multiple
228 acute studies Wilk [72,93,97] demonstrated that the shorter duration tempo e.g., faster movement produced the
229 highest 1RM value for bench press. These differences were notably found in movements of <2 s eccentric phases
230 compared to 4 and 6s [72] and 5, 8 and 10s [97] (Table 3). These findings reinforce the effects on concentric
231 performance such as an improved SSC response, use of stored elastic energy [23,78,107,108], and faster eccentric
232 velocities can result in the recruitment of higher threshold motor units [3,109]. Future research needs to determine
233 if performing eccentric phases <2 s has any greater effects on 1RM performance.

234 **4.3 Eccentric Tempo Chronic Response**

235 During chronic based studies to determine if 1RM increased utilising specific eccentric phase durations across the
236 same loads, a <2 s eccentric durations significantly increased back squat 1RM when compared to 4 s eccentric
237 durations which did not [68,98] (Table 5). However, findings on eccentric tempos are equivocal with Mike [97]
238 finding no difference in 1RM values between 2, 4 and 6 s eccentric phase (Table 5). Further, Kojić [70] found
239 that 4 s eccentric duration had a greater 1RM increases than 1 s eccentric tempos (23.5% vs 11.6%, respectively),
240 though both groups demonstrated significant strength increases. One explanation for this result is that the 4 s
241 group would have undergone significantly greater time under tension (TUT) and the 1 s group had demonstrated
242 significantly greater muscle volume increases which led to the concomitant increase in strength. Ünlü [60] found
243 no difference between 30 and 180°·s⁻¹ isotonic leg extension during eccentric actions. Douglas [43] found the
244 largest effect size during the 3 s AEL method to improve relative 1RM back squat, whereas, 3 s traditional
245 decreased scores and 1 s traditional slightly improved performance (Table 5).

246 The aforementioned chronic studies' findings indicate no clear duration which achieves a greater 1RM
247 enhancement (Table 5). Findings from Kojić [70] could speculate the 4 s group had greater TUT and hypertrophic
248 gain post study, which may have factored why there was an increase in 1RM. Furthermore, during the study by

249 Douglas [43] results indicated the 3 s AEL method achieved the greater improvement when compared to the other
250 methods. Though, it could be speculated to be an increase in motor learning as each training programme lasted
251 for 4 weeks. The acute findings of utilizing a <2 s eccentric tempo indicate an increase in peak and mean power
252 per repetition and 1RM increase. Though long-term training effects do not yield the same benefits of the acute
253 findings.

254 **4.4 Work and Time Under Tension**

255 TUT is the amount of time a muscle is held under tension or strain during an exercise set and therefore is the
256 product of duration in seconds in relation to how many repetitions are completed. When performing reduced
257 tempo durations compared with longer durations the amount of TUT will be less. However, performing a reduced
258 duration (faster tempo) enables performance of more repetitions [64]. The methodology behind this is, though
259 participants complete repetitions in a faster duration, they are able to complete more volume, subsequently,
260 increasing TUT [63,64,70,105]. This is solidified within the acute studies of [65,91] (Table 3), chronic study of
261 Kojić [70] (Table 5) and further support by Pryor [90] which indicated a higher total number of repetitions during
262 a 1 s eccentric tempo but did not show TUT results.

263 An important finding our review discovered is a number of studies viewed two different velocity, torque, or
264 duration values against one another. However, both training methods completed the same volume [65,67,94]
265 (Table 5 and 6). One could argue these comparative studies did not fulfil a similar protocol. Though, the same
266 volume is completed, metrics such as impulse and TUT will differ as the duration of the exercise is different. This
267 identifies that the same displacement was complete but with a reduced time and thus achieving a greater velocity,
268 resulting in a greater impulse. This is likely to factor results such as TUT, blood lactate (BL) and therefore,
269 hypertrophic response and strength improvements during chronic studies [110,111].

270 **4.5 Eccentric Execution Effects on Hypertrophy**

271 Eccentric training has been shown to increase hypertrophic gains greater than concentric training only
272 [12,14,105,112] via an increase in protein synthesis and IGF-1 mRNA expression as well as increases in
273 testosterone and growth hormone [105,110,113]. During acute tempo studies (Table 3) Wilk [65] identified that
274 a 6 s eccentric phase created larger TUT, BL, creatine kinase (CK) and serum testosterone compared to 2 s. Calixto
275 [67] had participants perform back squat with either 3 s or 0.5 s eccentric phase durations with the 3 s group
276 demonstrating significantly higher peak BL, 3, 6, 9, 15 and 20 mins post exercise BL and post 15 min growth
277 hormone compared to the 0.5 s. Martin-Costa [94] found a 4 s eccentric phase created a larger accumulation of

278 BL post 1 minute after 3 sets. Though the aforementioned studies [65,67,94] (Table 3 and 4) show metrics such
279 as testosterone and BL are enhanced during the longer durations tempos, caution must be taken when analysing
280 these results. As they all performed the same amount of volume with the same mass during the studies, thus the
281 tempos that are larger in duration will likely achieve a greater amount of TUT increasing BL which is correlated
282 with an increase in growth hormone response [110,111].

283 Shepstone [53] used a IKD protocol and highlighted moderate z-band disruption was greater for $210^{\circ}\cdot s^{-1}$ compared
284 to $20^{\circ}\cdot s^{-1}$, though extreme z-band disruption was not significant between groups, z-line streaming suggests
285 myofibrillar remodelling which leads to increase in muscle size [105] (Table 4). Nogueira [114] found no
286 significant difference between $20^{\circ}\cdot s^{-1}$ and $210^{\circ}\cdot s^{-1}$ for mechano-growth factor, total Akt, total mTOR and P70^{s6k1}.
287 However, both groups significantly improved from pre to immediately after and 2 hours post exercise for phospho-
288 Akt and P70^{s6k1} phosphorylation and both groups significantly improved phosphor-mTOR immediately after.
289 Though, $20^{\circ}\cdot s^{-1}$ showed that total impulse was greater (Table 4). Physiological occurrences such as MTOR and
290 Akt are indicative of anabolic signalling pathway, however, the limited results presented in the current review
291 identify no clear distinction between the training angular velocity values.

292 Stasinaki [68] found no difference between <1 s (as fast as possible under control) and 4 s eccentric only back
293 squat for both vastus lateralis muscle thickness and fascicle angle (Table 5). However, the <1 s group showed
294 significant differences compared to the 4 s group for fascicle length. Douglas [115] concluded there was no clear
295 difference between 1 s and 3 s (eccentric phase) traditional and AEL methods for muscle thickness and fascicle
296 angle and length (Table 5). It is important to note that both programmes only lasted for 4-weeks which may have
297 indicated no increase in muscle mass due to the physiological timelines needed to show muscle mass alterations
298 is typically longer than 4 weeks. Shibata [98] had increases in cross-sectional area (CSA) but no difference
299 between groups when comparing 2s eccentric and concentric and 4s eccentric 2s concentric during back squat to
300 failure. Kojić [70] highlighted no difference in bicep brachii muscle mass gain for 1s and 4s eccentric despite the
301 4s group achieving significantly greater eccentric TUT, though, the 1s group performed significantly greater
302 concentric TUT which has been shown to increase CSA [67,110] (Table 5). In conjunction with this, Fisher's
303 [116] review commented that there is no significant correlation between repetition duration and muscular
304 hypertrophy. This is further supported by Schoenfeld [117] who states there is no distinct difference between 0.8-
305 8s of total repetition duration with respect to muscle hypertrophy and eccentric duration should range from 2–4s.
306 Krzysztofik [63] further adds more research is needed [117] though eccentric phase durations of <2s show promise
307 for increasing muscle mass.

308 The current review further identifies IKD methods of $180\text{-}210^\circ\cdot\text{s}^{-1}$ and $20\text{-}30^\circ\cdot\text{s}^{-1}$ have been compared to
309 determine if increased angular velocity values create significant muscle mass difference (Table 6). The studies by
310 Farthing and Chilibeck [55,56] found no difference in increases for proximal, distal and mid site muscle thickness
311 (Table 6). However, one study [55] classified $180^\circ\cdot\text{s}^{-1}$ as a greater method to attain larger muscle thickness gains
312 due to a greater increase when compared to the $30^\circ\cdot\text{s}^{-1}$ and commented that greater eccentric angular velocities
313 lead to more muscle damage and therefore, greater hypertrophy (Table 6). Furthermore the isotonic study by Ünlü
314 [60] showed no difference in muscle mass between $180^\circ\cdot\text{s}^{-1}$ and $30^\circ\cdot\text{s}^{-1}$ for quadriceps muscle volume despite
315 each achieving 8–10s and 30–40s of TUT, respectively.

316 The study by Paddon-Jones [58] identified that $180^\circ\cdot\text{s}^{-1}$ significantly increased pre to post type IIb muscle fibre
317 values, but significantly decreased type I fibres (Table 6). The $30^\circ\cdot\text{s}^{-1}$ training procedure caused no significant
318 changes in any muscle fibres assessed and there was no difference between the 180 and $30^\circ\cdot\text{s}^{-1}$ post study.
319 Shepstone [53] (part 1) identified no difference between $210^\circ\cdot\text{s}^{-1}$ and $20^\circ\cdot\text{s}^{-1}$ eccentric IKD for type Ia and IIa fibre
320 %, though, type IIx significantly decreased for both groups from pre to post. The study did identify a significant
321 increase between the two methods for type IIa muscle fibre area % and myosin heavy chains. Additionally, both
322 groups significantly improved from pre to post testing for myosin heavy chains type IIa fibre and both significantly
323 decreased type IIx fibre % and myosin heavy chain (Table 6).

324 The review by Schoenfeld [105] further implies a distinct difference between isokinetic and isoinertial exercises
325 as during isokinetic there is a constant resistance and is not gravity dependent. Muscle tension has been categorised
326 as a key determinant for hypertrophy [105,118]. There will be an increase in muscular tension during faster
327 eccentric actions as both force and velocity are enhanced, therefore, muscular tension increases [14]. Schoenfeld
328 [117] comments that isokinetic training should be performed fast if the goal is to increase muscle mass, however,
329 the results of our review does not show clear differences between using the measures during eccentric actions.

330 Research seeks to optimise training to enable the greatest result with the available resources to practitioners to
331 elicit greater muscle mass, though, one could view that no difference between methods is a positive. First, this
332 means methods are interchangeable. However, as there is no distinct difference between methods, $<2\text{s}$ eccentric
333 tempos and $>180^\circ\cdot\text{s}^{-1}$ IKD can also increase other factors such as strength, power, and SSC function, it could be
334 argued that this method is superior because of its capacity to increase other aspects of performance. More research
335 is needed to investigate the effects of $<2\text{s}$ eccentric phases to determine how these impact muscle mass gains when
336 volume is not matched with a tempo that is greater than a 2s eccentric phase.

337 **4.6 Eccentric Velocity: What is Fast?**

338 Categorising fast and slow movements is a construct that researchers have based on movements where
339 comparisons are made between multiple specific velocity, torque or duration values against an action that is either
340 faster or slower than its counterpart. However, classification of the terms fast and slow can be misleading [45–
341 47]. Velocity-based training provides direct velocity ranges such as strength-speed ($0.75 - 1.00 \text{ m}\cdot\text{s}^{-1}$) and speed-
342 strength ($1.00 - 1.3 \text{ m}\cdot\text{s}^{-1}$ [13,37], avoiding use of the terms slow and fast. Table 7 demonstrates disparity within
343 the literature as to what is classified as fast (ranging from $180\text{-}240^\circ\cdot\text{s}^{-1}$) and slow ($20\text{-}60^\circ\cdot\text{s}^{-1}$) via Isokinetic
344 Dynamometry (IKD). Further the categorisation of fast and slow is an arbitrary process as the terms refer to no
345 set parameters [45,47]. Without such parameters, simply deeming one as fast due to its increased velocity/torque
346 is a comparative measure and does not necessarily follow that the other values may be considered slow. This
347 interpretation is supported by Cronin [45] who comments that utilising IKD at $60^\circ\cdot\text{s}^{-1}$ and $240\text{-}300^\circ\cdot\text{s}^{-1}$ (slow and
348 fast respectively) are classified to these terms simply due to the constraints of the equipment [13,16,48].

349

350

**** INSERT TABLE 7 HERE ****

351

352 To devise training programmes around velocity of movement a coach will typically manipulate the tempo of the
353 different phases of an exercise. Authors highlight that movement tempos should be performed for different
354 durations, classing these as fast and slow [14,61]. Reviews vary on the temporal definition of fast and slow tempos
355 with some suggesting that a 1 s eccentric and 1 s concentric is fast [14,61,62]. However, Krzysztofik's review
356 [63] defined fast as $<2 \text{ s}$ for the eccentric phase whereas [64] determined a temporal range of 2-9 s as fast for a
357 whole repetition, they explained durations $<2 \text{ s}$ could be termed as explosive or moving as fast as possible (Table
358 8). These disparities within the literature confuse how movements should be defined and performed to attain
359 desired results. Further complications in utilising phase duration to determine actions are linked to individual
360 anthropometrics. If two participants with different lever lengths complete a movement for the same duration, the
361 one with the greater range of motion would achieve a higher velocity, potentially achieving a different stimulus.

362

363

**** INSERT TABLE 8 HERE ****

364

365 Ambiguities in the calculations of movement velocities such as no consideration of participant's anthropometrics
366 coupled with the use of the terms 'fast' and 'slow' should be altered to improve accuracy in the literature. For
367 example, when comparing training methods such as tempo training and IKD the terms of fast and slow become
368 redundant, as they can only be termed in their specific training method and cannot be compared to other training
369 methods. Therefore, within future research, specific values should be utilised alone. If authors wish to add tempo
370 terms to intended durations they are advised to view research by Wilk et al [64] who incorporates exercise tempos
371 of fast (2-4.9 s), medium (5-9.9 s) and slow (10-14.9 s) terms. With the addition of X providing maximal velocity
372 the load can be lifted and V which determines an individual's preferred velocity during an exercise (tempo
373 example X/1/2/1). If all future studies use the same terms of tempo training this would avoid confusion during
374 their implementation.

375 **4.7 Accelerated Eccentrics: The Resurrection**

376 Various terms are utilised to describe an intended accelerated eccentric motion: fast eccentric, dynamic eccentrics,
377 antagonistic facilitation specialised method (AFSM), overspeed eccentrics, accentuated eccentric loading, and
378 accelerated eccentrics. Having no agreed operational definition limits both research and practical implementation
379 as various terms are used in accordance with similar or the same intended training method. Verkoshansky and Siff
380 [13] comment on implementing "accelerated powermetrics" motions during training. They termed this
381 "accelerated eccentrics" which is sub-categorised into two groups. The first is passively accelerated eccentrics
382 where either resistance bands are incorporated, or an external partner pushes the barbell (a method of an inverse
383 forced repetition). The aim is to enable the bar to accelerate downwards and when the desired distance is
384 completed the athlete will decelerate the bar and complete the concentric phase. Actively accelerated eccentrics
385 are when an individual will perform a drop catch where the intention is to begin with limbs at extension and then
386 to drop and catch the bar towards the end of the eccentric phase. Another form of actively accelerated eccentrics
387 is to pull the bar down as fast as possible. Therefore, the term accelerated eccentrics would be the most applicable
388 to utilise as it focuses on maximising the acceleration of the eccentric phase and incorporates an agreed definition
389 to the training method reducing confusion among practitioners and researchers.

390 The term overspeed eccentrics originates from Louie Simmons of Westside Barbell Club who used resistance
391 bands to increase eccentric acceleration and provide accommodating resistance. However, overspeed is a term
392 widely used in coaching and is given to an exercise that achieves a greater velocity than is possible during normal
393 isoinertial movements and can cause confusion as the term overspeed is used to define two differing types of

394 training. Overspeed eccentrics as a term should not be used in accordance with assisted training where an athlete's
395 load is decreased by attaching bands to a harness [73] or by pulling on a band during a jump [74]. The term
396 overspeed eccentrics can still be utilised by coaches, though, caution must be applied for the term not to be used
397 for assisted training to ensure the integrity of the term remains clear throughout literature. To the authors
398 knowledge AFSM has not yet be referenced in the peer reviewed literature, the method originates from Triphasic
399 training programme [75].

400 Resistance bands have been used during AEL to provide resistance during the eccentric phase of a
401 countermovement jump and drop jump [76,77]. Though this method would accelerate an individual's mass faster
402 during the eccentric phase than weighted AEL [51] with the aim to accelerate an athlete incorporating the velocity
403 measure in the force-velocity curve [37]. Therefore, it is recommended that if bands are utilised instead of
404 increasing mass during AEL then the term of accelerated eccentric loading (ACEL) should be employed as this
405 give further credence to the training method. This can be seen in the band release method [40].

406 **4.8 Accelerated Eccentric Training Methods: Current Evidence**

407 When interpreting values from IKD, it is difficult to compare it with isoinertial movement values, which are
408 gravity dependent [52,78]. During IKD there is an initial acceleration phase [13] followed by a period of constant
409 torque, the purpose being to resist force during eccentric actions [79]. During isoinertial methods such as free
410 weights, acceleration due to gravity is the only external force acting on a mass. Therefore, when performing high
411 velocity movements during an unweighing phase (when individuals do not attempt to decelerate a movement)
412 there is no mechanical tension being applied to the mass [78]. This is how IKD and isoinertial methods differ [80],
413 however, once athletes decelerate the mass (braking phase) eccentric muscle lengthening occurs [78].
414 Additionally, during accelerated eccentric motions mass travels at a greater velocity. If two objects are the same
415 mass and travel the same displacement, though if one has a greater acceleration, then momentum will increase,
416 subsequently, a greater impulse (impulse = force \times Δ time) is required to stop the mass in motion [51]. This is
417 evidenced by Van den Tillaar [81], where participants performed back squats with different eccentric execution
418 velocities, they identified faster eccentric actions achieved the greatest force as the two faster velocities achieved
419 a greater peak force during the lowest displacement (bottom) of a back squat [7].

420 A recent survey exploring eccentric training methods demonstrated that IKD is rarely implemented within training
421 [82,83] as it has little practical application within a strength and conditioning environment and that very few have
422 access to [80,84]. Training should replicate sporting movements [52], supporting the theory of dynamic

423 correspondence [13,75,85,86], reinforcing that IKD is limited to angular velocity values that are not compatible
424 with sporting motions [45]. Interestingly, the aforementioned studies highlighted tempo training is frequently
425 employed in practice [82,83,87]. Thus, suggesting that research must provide more practical based methods such
426 as tempo training to enable a greater impact on larger populations. Further, Paddon-Jones [58] eluded that fast
427 and slow are incorrect terms, and that methods should specify training velocities, as specific values create different
428 results. This implies that additional methods such as eccentric cueing or resistance banded exercises need to be
429 understood to enable coaches to employ accelerated eccentrics more effectively.

430 **5.0 Conclusion**

431 We as the authors, recommend when using increased velocity eccentric actions, the term of accelerated eccentrics
432 should be used and accelerated eccentric loading (ACEL) should be employed when the external accelerating
433 stimulus is released at the end of the eccentric phase. This creates a distinct difference between increasing mass
434 (AEL) or velocity (ACEL). When employing IKD the terms of fast and slow should be avoided, instead specific
435 angular velocity values should be reported. During tempo training the term of fast or <1 s tempo should be
436 substituted with “as fast as possible” furthermore, during tempo training specific timeframes should be employed
437 instead of the previous fast and slow classification (Table 1).

438 If IKD is utilised, angular velocities of $>180^{\circ}\text{s}^{-1}$ should be used over $<60^{\circ}\text{s}^{-1}$ as results imply greater strength
439 gains. There is no clear difference between $<60^{\circ}\text{s}^{-1}$ and $>180^{\circ}\text{s}^{-1}$ for hypertrophic increase. Tempo methods of
440 $<2\text{s}$ are superior to create greater 1RM scores and higher mean and peak concentric velocity and power. Slower
441 tempos ($>2\text{s}$) result in greater accumulation of BL, growth hormone and testosterone however, this only occurs
442 when the volume and intensity is matched. Research needs to review shorter durations with greater volume than
443 longer ones to determine if one is superior in achieving greater muscle mass. There is no clear difference between
444 $<2\text{ s}$ and $>4\text{s}$ for increases in muscle mass. This review highlights the need for more research into training with
445 various eccentric speeds with particular focus on areas such as accelerated eccentrics and $<1\text{ s}$ tempos to determine
446 adaptations from these methods.

Reference List

1. Lindstedt SL, LaStayo PC, Reich TE. When active muscles lengthen: Properties and consequences of eccentric contractions. *News Physiol Sci.* 2001;16:256–61.
2. Harris-Love MO, Seamon BA, Gonzales TI, Hernandez HJ, Pennington D, Hoover BM. Eccentric exercise program design: A periodization model for rehabilitation applications. *Front Physiol.* 2017;8:1–16.
3. Douglas J, Pearson S, Ross A, McGuigan M. Eccentric exercise: physiological characteristics and acute responses. *Sport Med.* Springer International Publishing; 2017;47:663–75.
4. Verkhoshansky Y. *Fundamentals of special strength-training in sport.* 1986.
5. Isner-Horobeti ME, Dufour SP, Vautravers P, Geny B, Coudeyre E, Richard R. Eccentric exercise training: modalities, applications and perspectives. *Sport Med.* 2013;43:483–512.
6. Katz BYB. The relation between force and speed in muscular contraction. *J Physiol.* 1939;96:45–64.
7. Mcneill C, Beaven CM, McMaster DT, Gill N. Eccentric force-velocity characteristics during a novel squat protocol in trained rugby union athletes — pilot study. *J Funct Morphol Kinesiol.* 2021;6:1–10.
8. Tobin DP. Advanced strength and power training for the elite athlete. *Strength Cond J.* 2014;36:59–65.
9. Wagle JP, Taber CB, Cunanan AJ, Bingham GE, Carroll KM, DeWeese BH, et al. Accentuated eccentric loading for training and performance: a review. *Sport Med.* Springer International Publishing; 2017;47:2473–95.
10. Wagle J., Cunanan A., Carroll K., Sams M., Wetmore A, Bingham G., et al. Accentuated eccentric loading and cluster set configurations in the back squat: a kinetic and kinematic analysis. *J Strength Cond Res.* 2018;00:1–8.
11. Watkins PH. Augmented eccentric loading: theoretical and practical applications. *UKSCA J [Internet].* 2010;3–11. Available from: <http://www.scopus.com/inward/record.url?eid=2-s2.0-84896553908&partnerID=40&md5=f39fed844b11d966ba71c100bee3d146>
12. Douglas J. The role of eccentric muscle function and training in athletic performance. Doctoral dissertation. Auckland University of Technology; 2018.
13. Vaerhoshanky, Y & Siff M. *Supertraining.* Rome Italy: Verkhoshanky SSTM; 2009.

14. Suchomel TJ, Wagle JP, Douglas J, Taber CB, Harden M, Haff GG, et al. Implementing eccentric resistance training — part 1 : a brief review of existing methods. *J Funct Morphol Kinesiol*. 2019;4:1–25.
15. Suchomel TJ, Nimphius S, Bellon CR, Stone MH. The importance of muscular strength: training considerations. *Sport Med* [Internet]. Springer International Publishing; 2018;48:765–85. Available from: <https://doi.org/10.1007/s40279-018-0862-z>
16. Zatsiorsky MV, Kraemer WJ. Science and practice of strength training [Internet]. 2006. p. 264. Available from: <http://books.google.com/books?hl=en&lr=&id=QWSn4iKgNo8C&oi=fnd&pg=PR8&dq=Science+and+Practice+of+Strength+Training&ots=v46bUiOLDs&sig=qduV5QRbeak7oxUarrpS0XNBixY%5Cnhttps://books.google.com/books?id=QWSn4iKgNo8C&pgis=1>
17. McNeill C, Beaven CM, McMaster DT, Gill N. Eccentric training interventions and team sport athletes. *J Funct Morphol Kinesiol*. 2019;4:67.
18. Pereira MIR, Gomes PSC. Movement velocity in resistance training. *Sport Med*. 2003;33:427–38.
19. Wilkie DR. The relation between force and velocity in human muscle. *J Physiol*. 1949;110:249–80.
20. Haeufle DFB, Günther M, Bayer A, Schmitt S. Hill-type muscle model with serial damping and eccentric force-velocity relation. *J Biomech*. 2014;47:1531–6.
21. Brughelli, M., Cronin J. Altering the length-tension relationship with eccentric exercise. *Sport Med*. 2007;37:807–26.
22. Herzog W. The role of titin in eccentric muscle contraction. *J Exp Biol*. 2014;217:2825–33.
23. Duncan PW, Chandler JM, Cavanaugh DK, Johnson KR, Buehler AG. Mode and speed specificity of eccentric and concentric exercise training. *J Orthop Sports Phys Ther*. 1989;11:70–5.
24. Hildre T. The effects of velocity on force production of the elbow flexors during eccentric isokinetic muscle contraction. 1996.
25. Cress NM, Peters KS, Chandler JM. Eccentric and concentric force-velocity relationships of the quadriceps femoris muscle. *J Orthop Sports Phys Ther*. 1992;16:82–6.
26. Cabri JMH. Isokinetic strength aspects of human joints and muscles. *Crit Rev Biomed Eng*. 1991;19:231–

59.

27. Chapman D, Newton M, Nosaka K. Eccentric torque-velocity relationship of the elbow flexors. *Isokinet Exerc Sci.* 2005;13:139–45.

28. Yeadon MR, King MA, Wilson C. Modelling the maximum voluntary joint torque/angular velocity relationship in human movement. *J Biomech.* 2006;39:476–82.

29. Cowell JF, Cronin J, Brughelli M. Eccentric muscle actions and how the strength and conditioning specialist might use them for a variety of purposes. *Strength Cond J.* 2012;34:33–48.

30. Lee EJ, Joumaa V, Herzog W. New insights into the passive force enhancement in skeletal muscles. *J Biomech.* 2007;40:719–27.

31. Herzog W, Schappacher G, DuVall M, Leonard T., Hezong J. Residual force enhancement following eccentric contractions: a new mechanism involving titin. *Physiology.* 2016;31:300–12.

32. Powers K, Schappacher-Tilp G, Jinha A, Leonard T, Nishikawa K, Herzog W. Titin force is enhanced in actively stretched skeletal muscle. *J Exp Biol.* 2014;217:3629–36.

33. Leonard TR, Herzog W. Regulation of muscle force in the absence of actin-myosin-based cross-bridge interaction. *Am J Physiol - Cell Physiol.* 2010;299:14–20.

34. Campos DB, Ferreira IC, Souza MA, Amorim M, Intelangelo L, Silveira-Nunes G, et al. Acceleration profiles and the isoinertial squatting exercise: is there a direct effect on concentric–eccentric force, power, and neuromuscular efficiency? *J Sport Rehabil.* 2020;30:646–52.

35. Schilling BK, Falvo MJ, Chiu LZF. Force-velocity, impulse-momentum relationships: Implications for efficacy of purposefully slow resistance training. *J Sport Sci Med.* 2008;7:299–304.

36. Young W, Bilby G. The effect of voluntary effort to influence speed of contraction on strength, muscular power, and hypertrophy development. *J Strength Cond Res.* 1993;7:172–8.

37. Zatsiorsky V. *Biomechanics in sport: performance enhancement and injury prevention.* Oxford: Blackwell Science Ltd. 2000.

38. Verkhoshansky Y V. *Shock method.* Rome Italy: Verkhoshanky SSTM; 2018.

39. Verkhoshansky, Y. Verkhoshansky N. *Special strength training manual for coaches.* Rome Italy:

Verkhoshanky SSTM; 2011.

40. Handford MJ, Rivera FM, Maroto-Izquierdo S HJ. Plyo-accentuated eccentric loading methods to enhance lower limb muscle power. *Strength Cond J.* 2021;43:54–64.
41. Wagle J, Taber C, Carroll K, Cunanan A, Sams M, Wetmore A, et al. Repetition-to-repetition differences using cluster and accentuated eccentric loading in the back squat. *Sports [Internet].* 2018;6:59. Available from: <http://www.mdpi.com/2075-4663/6/3/59>
42. Lates AD, Greer BK, Wagle JP, Taber CB. Accentuated eccentric loading and cluster set configurations in the bench press. *J Strength Cond Res.* 2020;Publish Ah:1–5.
43. Douglas J, Pearson S, Ross A, McGugan M. Effects Of Accentuated Eccentric Loading On Muscle Properties, Strength, Power, And Speed In Resistance-Trained Rugby Players. *J Strength Cond Res.* 2018;32:2750–61.
44. Harden M, Worf A, Russell M, Hicks K., French D, Howatson G. An evaluation of supramaximally loaded eccentric leg press exercis. *J Strength Cond Res.* 2018;32:2708–14.
45. Cronin JB, McNair PJ, Marshall RN. Is velocity-specific strength training important in improving functional performance? *J Sports Med Phys Fitness.* 2002;42:267–73.
46. Hedrick A. Literature review: high speed resistance training. *Natl Strength Cond Assoc J.* 1993;15:22–30.
47. Cronin J, McNair PJ, Marshall RN. Velocity specificity, combination training and sport specific tasks. *J Sci Med Sport.* 2001;4:168–78.
48. Andrade MDS, Fleury AM, de Lira CAB, Dubas JP, da Silva AC. Profile of isokinetic eccentric-to-concentric strength ratios of shoulder rotator muscles in elite female team handball players. *J Sports Sci.* 2010;28:743–9.
49. Iossifidou A, Baltzopoulos V, Giakas G. Isokinetic knee extension and vertical jumping : Are they related ? *2005;23:1121–7.*
50. Chapman D, Newton M, Sacco P, Nosaka K. Greater muscle damage induced by fast versus slow velocity eccentric exercise. *Int J Sports Med.* 2006;27:591–8.
51. Frost DM, Cronin J, Newton RU. A biomechanical evaluation of resistance: fundamental concepts for

- training and sports performance. *Sport Med.* 2010;40:303–26.
52. Kellis E, Baltzopoulos V. Isokinetic eccentric exercise. *Sport Med.* 1995;19:202–22.
53. Shepstone TN, Tang JE, Dallaire S, Schuenke MD, Staron RS, Phillips SM. Short-term high- vs. low-velocity isokinetic lengthening training results in greater hypertrophy of the elbow flexors in young men. *J Appl Physiol.* 2005;98:1768–76.
54. Roschel H, Ugrinowitsch C, Barroso R, Batista MAB, Souza EO, Aoki MS, et al. Effect of eccentric exercise velocity on akt/mTOR/p70s6k signaling in human skeletal muscle. *Appl Physiol Nutr Metab.* 2011;36:283–90.
55. Farthing JP, Chilibeck PD. The effects of eccentric and concentric training at different velocities on muscle hypertrophy. *Eur J Appl Physiol.* 2003;89:578–86.
56. Farthing JP, Chilibeck PD. The effect of eccentric training at different velocities on cross-education. *Eur J Appl Physiol.* 2003;89:570–7.
57. Paddon-Jones, D., Keech, A., Lonergan, A., & Abernethy P. Differential expression of muscle damage in humans following acute fast and slow velocity eccentric exercise. *J Sci Med Sport.* 2005;8:255–63.
58. Paddon-Jones, D., Leveritt, M., Lonergan, A., & Abernethy P. Adaptation to chronic eccentric exercise in humans: the influence of contraction velocity. *Eur J Appl Physiol.* 2001;85:466–71.
59. Aagaard P, Simonsen EB, Andersen JL, Magnusson SP, Halkjær-Kristensen J, Dyhre-Poulsen P. Neural inhibition during maximal eccentric and concentric quadriceps contraction: effects of resistance training. *J Appl Physiol.* 2000;89:2249–57.
60. Ünlü G, Çevikol C, Melekoğlu T. Comparison of the effects of eccentric, concentric, and eccentric-concentric isotonic resistance training at two velocities on strength and muscle hypertrophy. *J strength Cond Res.* 2020;34:337–44.
61. Davies TB, Kuang K, Orr R, Halaki M, Hackett D. Effect of movement velocity during resistance training on dynamic muscular strength: a systematic review and meta-analysis. *Sport Med. Springer International Publishing;* 2017;47:1603–17.
62. Ratamess, NA, Alvar, BA, Evetoch, TE, Housh, TJ, Ben Kibler, W, Kraemer, WJ, Triplett N. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc.* 2009;41:687–708.

63. Krzysztofik M, Wilk M, Wojdała G, Gołaś A. Maximizing muscle hypertrophy: a systematic review of advanced resistance training techniques and methods. *Int J Environ Res Public Health*. 2019;16.
64. Wilk M, Tufano JJ, Zajac A. The influence of movement tempo on acute neuromuscular, hormonal, and mechanical responses to resistance exercise—a mini review. *J strength Cond Res*. 2020;34:2369–83.
65. Wilk M, Stastny P, Golas A, Nawrocka M, Jelen K, Zajac A, et al. Physiological responses to different neuromuscular movement task during eccentric bench press. *Neuroendocrinol Lett*. 2018;39:26–32.
66. Wilk M, Golas A, Krzysztofik M, Nawrocka M, Zajac A. The effects of eccentric cadence on power and velocity of the bar during the concentric phase of the bench press movement. *J Sport Sci Med*. 2019;18:191–7.
67. Calixto RD, Verlengia R, Crisp AH, Carvalho TB, Crepaldi MD, Pereira AA, et al. Acute effects of movement velocity on blood lactate and growth hormone responses after eccentric bench press exercise in resistance-trained men. *Biol Sport*. 2014;31:289–94.
68. Stasinaki A-N, Zaras N, Methenitis S, Bogdanis G, Terzis G. Rate of Force Development and Muscle Architecture after Fast and Slow Velocity Eccentric Training. *Sports*. 2019;7:1–12.
69. Pereira PE, Motoyama Y, Tanaka K. Resistance training with slow speed of movement is better for hypertrophy and muscle strength gains than fast speed of movement. *Int J Appl Exerc Physiol*. 2016;13:466.
70. Kojić, F., Ranisavljev, I., Čosić, D., Popović, D., Stojiljković, S., & Ilić V. Effects of resistance training on hypertrophy, strength and tensiomyography parameters of elbow flexors: role of eccentric phase duration. *Biol Sport*. 2021;38:587–94.
71. Wilk M, Gepfert M, Krzysztofik M, Golas A, Mostowik A, Maszczyk A, et al. The influence of grip width on training volume during the bench press with different movement tempos. *J Hum Kinet*. 2019;68:49–57.
72. Lopes CR, Crisp AH, Rodrigues AL, Teixeira AG, Da Mota GR, Verlengia R. Fast contraction velocity in resistance exercise induces greater total volume load lifted and muscle strength loss in resistance-trained men. *Rev Andaluza Med del Deport*. 2012;5:123–6.
73. Argus CK, Gill ND, Keogh JW, Blazevich AJ, Hopkins WG. Kinetic and training comparisons between assisted, resisted, and free countermovement jumps. *J Strength Cond Res*. 2011;25:2219–27.
74. Stien N, Strate M, Andersen V, Saeterbakken AH. Effects of Overspeed or Overload Plyometric Training on Jump Height and Lifting Velocity. *Sport Med Int open*. 2020;4:32–8.

75. Dietz. C PB. Triphasic training. 2012.
76. Aboodarda S., Byrne J., Samson M, Wilson B., Okhtar A., Behm D. Does performing drop jumps with additional eccentric loading improve jump performance. *J Strength Cond Res.* 2014;28:2314–23.
77. Aboodarda S., Yusof A, Osman NA., Thompson M., Mokhtar A. Enhanced performance with elastic resistance during the eccentric phase of a countermovement jump. *Int J Sport Physiol Perform.* 2013;8:181–7.
78. Guilhem G, Cornu C, Gue A. Neuromuscular and muscle-tendon system adaptations to isotonic and isokinetic eccentric exercise. *Ann Phys Rehabil Med.* 2010;53:319–41.
79. Zatsiorsky, VM, & Prilutsky B. Biomechanics of skeletal muscles. *Hum Kinet.* 2012. p. 1–519.
80. Meylan C, Cronin J, Nosaka K. Isoinertial assessment of eccentric muscular strength. *Strength Cond J.* 2008;30:56–64.
81. Van den Tillaar R. Effect of descent velocity upon muscle activation and performance in two-legged free weight back squats. *Sports.* 2019;7:15.
82. McNeill C, Beaven CM, McMaster DT, Gill N. Survey of eccentric-based strength and conditioning practices in sport. *J strength Cond Res.* 2020;34:2769–75.
83. Harden. M, Bruce. C, Wolf. A, Hicks. K.M HG. Exploring the practical knowledge of eccentric resistance training in high- performance strength and conditioning practitioners. *Int J Sport Sci Coach.* 2019;15:1–12.
84. Coratella G, Milanese C, Schena F. Unilateral eccentric resistance training: a direct comparison between isokinetic and dynamic constant external resistance modalities. *Eur J Sport Sci.* 2015;15:720–6.
85. Verkhoshansky YU, Yessis M. *Special Strength Training: A Practical Manual for Coaches* [Internet]. 2006. p. 137. Available from: files/1721/Verkhoshansky and Verkhoshansky - 2011 - Special Strength Training Manual for Coaches.pdf
86. Suarez DG, Wagle JP, Cunanan AJ, Sausaman RW SM. Dynamic correspondence of resistance training to sport: a brief review. *Strength Cond J.* 2019;41:80–8.
87. Drury B, Clarke H, Moran J, Fernandes JFT, Henry G, Behm DG. Eccentric resistance training in youth : a survey of perceptions and current practices by strength and conditioning coaches. *J Funct Morphol Kinesiol.* 2021;6:1–14.

88. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health*. 1998;52:377–84.
89. Carzoli, J P, Sousa, C.A, Belcher, D.J, Helms, ER, Khamoui, AV, Whitehurst, M, Zourdos M. The effects of eccentric phase duration on concentric outcomes in the back squat and bench press in well-trained males. *J Sports Sci*. 2019;37:2676–84.
90. Pryor RR, Sforz GA, Kin DL. Optimizing power output by varying repetition tempo. *J Strength Cond Res*. 2011;25:3029–34.
91. Sampson JA, Donohoe A, Groeller H. Effect of concentric and eccentric velocity during heavy-load non-ballistic elbow flexion resistance exercise. *J Sci Med Sport*. 2014;17:306–11.
92. Wilk M, Golas A, Zmijewski P, Krzysztofik M, Filip A, Coso J Del, et al. The effects of the movement tempo on the one-repetition maximum bench press results. *J Hum Kinet*. 2020;72:151–9.
93. Wilk M, Gepfert M, Krzysztofik M, Mostowik A, Filip A, Hajduk G, et al. Impact of duration of eccentric movement in the one-repetition maximum test result in the bench press among women. *J Sport Sci Med*. 2020;19:317–22.
94. Martins-Costa HC, Diniz RCR, Lima FV, Machado SC, De Almeida RSV, De Andrade AGP, et al. Longer repetition duration increases muscle activation and blood lactate response in matched resistance training protocols. *Motriz Rev Educ Fis*. 2016;22:35–41.
95. Drury DG, Stuempfle KJ, Mason CW, Girman JC. The effects of isokinetic contraction velocity on concentric and eccentric strength of the biceps brachii. *J Strength Cond Res*. 2006;20:390–5.
96. Hortobágyi T, Katch FI. Eccentric and concentric torque-velocity relationships during arm flexion and extension - Influence of strength level. *Eur J Appl Physiol Occup Physiol*. 1990;60:395–401.
97. Mike JN, Cole N, Herrera C, Vandusseldorp T, Kravitz L, Kerksick CM. The effects of eccentric contraction duration on muscle strength, power production, vertical jump, and soreness. *J Strength Cond Res*. 2017;31:773–86.
98. Shibata.k, Takzawa. K, Noska. K MM. Effects of prolonging eccentric phase duration in parallel back-squat training to momentary failure on muscle cross-sectional area, squat one repetition maximum, and performance

- tests in university soccer players. *J Strength Cond Res.* 2018;35:668–74.
99. Y.Yamada NI. Differences in eccentric force-velocity characteristics between isotonic and isokinetic contractions. *J Sport Heal Sci.* 2018;121.
100. Katz BYB. The relation between force and speed in muscular contraction. *J Physiol.* 1939;96:45–64.
101. Roig M, O'Brien K, Kirk G, Murray R, McKinnon P, Shadgan B, et al. The effects of eccentric versus concentric resistance training on muscle strength and mass in healthy adults: A systematic review with meta-analysis. *Br J Sports Med.* 2009;43:556–68.
102. Douglas J, Pearson S, Ross A, McGuigan M. Chronic adaptations to eccentric training: a systematic review. *Sport Med.* Springer International Publishing; 2017;47:917–41.
103. Hody S, Croisier JL, Bury T, Rogister B, Leprince P. Eccentric muscle contractions: Risks and benefits. *Front Physiol.* 2019;10:1–18.
104. Ryan LM, Magidow PS, Duncan PTPW. Velocity-specific and mode-specific effects of eccentric isokinetic training of the hamstrings. *J Orthop Sport Phys Ther.* 1991;13:33–9.
105. Schoenfeld BJ. The mechanisms of muscle hypertrophy and their application to resistance training. *J Strength Cond Res.* 2010;24:2857–72.
106. Richards CT, Sawicki GS. Elastic recoil can either amplify or attenuate muscle-tendon power, depending on inertial vs. fluid dynamic loading. *J Theor Biol.* Elsevier; 2012;313:68–78.
107. Lindstedt SL, Reich TE, Keim P, LaStayo PC. Do muscles function as adaptable locomotor springs? *J Exp Biol.* 2002;205:2211–6.
108. Behm DG, Sale DG. Velocity specificity of resistance training. *Sport Med.* 1993;15:374–88.
109. Nardone, A., Romano, C., & Schieppati M. Selective recruitment of high-threshold human motor units during voluntary isotonic lengthening of active muscles. *J Physiol.* 1989;409:451–71.
110. Kraemer RR, Castracane VD. Endocrine alterations from concentric vs. eccentric muscle actions: a brief review. *Metabolism [Internet].* Elsevier Inc.; 2015;64:190–201. Available from: <http://dx.doi.org/10.1016/j.metabol.2014.10.024>
111. Godfrey RJ, Whyte G. The role of lactate in the exercise-induced growth hormone response : evidence

from myocardial disease the role of lactate in the exercise-induced human growth hormone response : evidence from myocardial disease. *Br J Sports Med.* 2005;43:521–5.

112. Schoenfeld BJ, Ogborn DI, Vigotsky AD, Franchi M V., Krieger JW. Hypertrophic effects of concentric vs. eccentric muscle actions: a systematic review and meta-analysis. *J Strength Cond Res.* 2017;31:2599–608.

113. Komi P V. *Strength and power in sport.* John Wiley Sons. 2008.

114. Nogueira FRD, Conceição MS, Vechin FC, Mendes EM, Rodrigues GFC, Fazolin MA, et al. The effect of eccentric contraction velocity on muscle damage: A review. *Isokinet Exerc Sci.* 2013;21:1–9.

115. Douglas J, Pearson S, Ross A, McGuigan M. Effects of accentuated eccentric loading on muscle properties, strength, power, and speed in resistance-trained rugby players. *J Strength Cond Res.* 2018;32:2750–61.

116. J. Fisher, J. Steele DS. Evidence-based resistance training recommendations for muscular hypertrophy. *Med Sport.* 2013;17:217–35.

117. Schoenfeld BJ, Ogborn DI, Krieger JW. Effect of repetition duration during resistance training on muscle hypertrophy: a systematic review and meta-analysis. *Sport Med.* 2015;45:577–85.

118. Suchomel TJ, Wagle JP, Douglas J, Taber CB, Harden M, Ha GG, et al. Implementing eccentric resistance training — part 2 : practical recommendations. *J Funct Morphol Kinesiol.* 2019;4:1–20.

Table 1: Current review terms and definitions of said terms.

Term	Acronym	Definition
Accelerated Eccentric	NA	The training method whereby, the aim is to perform the eccentric phase at an increased velocity, determined by eccentric phase execution strategies.
Overspeed Eccentric	NA	The addition of resistance bands by causing the subject to move faster on the eccentric phase producing more power for reversal strength.
Accelerated Eccentric Loading	ACEL	The training method where resistance bands are employed to accelerate the eccentric phase of an exercise and the bands are then released prior to the concentric phase of an exercise.
Accentuated Eccentric Loading	AEL	The training method whereby the eccentric phase of an eccentric concentric coupling exercise has an increased mass than when compared to the concentric phase.
Isokinetic Dynamometry	IKD	The training method whereby, an isokinetic device is utilised to complete an exercise.
Tempo Training	NA	The training method whereby, an allotted time duration is given in respect to the eccentric, isometric, concentric and isometric phases of an exercise.
Fast tempo	NA	Term should be avoided, however, if used it defines a tempo that is faster than its counterpart.
Slow Tempo	NA	Term should be avoided, however, if used it defines a tempo that is slower than its counterpart.
Isoinertial	NA	When mass is consistent during resistance-based training.

NA: Not applicable.

Table 2: Study eligibility criteria.

1.	Peer reviewed journal article displayed (full text).
2.	English language.
3.	Study published within the years of 1950–2021.
4.	Study was conducted on both male and female participants.
5.	Participants must have been healthy and have no injuries, disability, or illness.
6.	Participants were aged 18-50 years of age.
7.	Analysed the acute and/or chronic effects of eccentric training.
8.	Study must have viewed more than one eccentric: duration, speed, velocity, and torque values.
9.	Studies must have completed research via a comparative measure.
10.	Concentric performance if tested must have been the same across testing.

Table 3: Acute tempo studies.

Study	Population	Aim	Method	Findings
Van den Tillaar [81]	N = 11 healthy resistance trained males age = 24.0 ± 6.0 yrs, BM = 89.5 ± 21.5 kg, height = 184.0 ± 10.0 cm, TE = > 2.0 yrs, 4RM BS = 129.0 ± 23.0 kg.	To determine the kinematic results of different ECC TEMPOS on a 4RM BS.	Exercise: BS, intensity 4RM, ECC TEMPO: TEMPO 1: participant's normal decent velocity, TEMPO 2: decent at a faster velocity than TEMPO 1. TEMPO 3: decent slower than normal velocity. All velocities were subjected to the participants in the study. The descent velocity was on average $0.3 \text{ m}\cdot\text{s}^{-1}$ faster and slower in TEMPO 2 and 3 velocities compared to the TEMPO 1, respectively. VDOWN: Negative acceleration before deceleration. V0: Lowest displacement of squat. VMAX1: Peak acceleration prior to sticking point. VMIN: Sticking point. VMAX2: Second peak velocity score.	VDOWN: Significant increase in displacement, velocity, and duration for TEMPO 2 vs TEMPO 1 and 3. V0: Significant increase in displacement for TEMPO 2 vs TEMPO 1 and 3. Significant increase in peak force for TEMPO 2 and 1 vs TEMPO 3. VMAX1: TEMPO 2 and 1 vs TEMPO 3 had significantly greater duration and velocity. VMIN: No difference in TEMPOS for measures assessed. VMAX2: No difference in TEMPOS for measures assessed.
Wilk [66]	N = 33 male, age = 24.0 ± 4.2 yrs, BM = 77.3 ± 5.7 kg, BP 1RM = 107.4 ± 13.5 kg, TE: 2.2 ± 0.6 yrs.	The effects of 2 s and 6 s ECC cadence on concentric power and velocity.	Exercise: BP, volume: 3 x failure, intensity: 70% of 1RM, ECC tempos: ECC-2s: 2s ECC, X CON ECC-6s: 6s ECC, X CON	Increase in mean velocity ($\text{m}\cdot\text{s}^{-1}$) for ECC-2s vs ECC-6s: Set 1 (15.38%), Set 2 (19.23%), Set 3 (21.57%). Increase in peak velocity ($\text{m}\cdot\text{s}^{-1}$) for ECC-2s vs ECC-6s: Set 1 (14.49%), Set 2 (17.14%), Set 3 (19.12%). Increase in mean power (W) for ECC-2s vs ECC-6s: Set 1 (16.46%) Set 2 (20.00%), Set 3 (23.97%). Increase in peak power (W) for ECC-2s vs ECC-6s: Set 1 (18.71%), Set 2 (19.45%), Set 3 (26.64%). Significant increase for ECC-2s vs ECC-6s: groups for all sets and all metrics.
Carzoli [89]	N = 16 trained males age = 23.6 ± 2.8 yrs, height = 171.8 ± 7.5 cm, BM = 82 ± 12.2 kg, BS 1RM = 151.8 ± 49.6 kg,	To determine the effects of different ECC TEMPOS on average and peak power scores.	Exercise: BS and BP, intensity: 60 and 80% of 1RM during both exercises. ECC TEMPOS: Controlled by metronome:	Increase in BS mean CON velocity ($\text{m}\cdot\text{s}^{-1}$) 60% 1RM: TEMPO 1 vs TEMPO 2 (8.06%), and TEMPO 3 (10.04%), TEMPO 1 significantly greater than both.

	BP 1RM = 119.7 ± 26.2, TE = 7.0 ± 3.6 yrs.		TEMPO 1: 0.75 times participant's normative ECC mean squat at 60 and 80% 1RM for both BS and BP. TEMPO 2: Normative. TEMPO 3: 2.00 times participant's normative ECC mean squat at 60 and 80% 1RM for both BS and BP.	Increase in BS peak CON velocity (m·s ⁻¹) 60% of 1RM: TEMPO 1 vs TEMPO 2 (74.44%) and TEMPO 3 (9.21%), TEMPO 1 significantly greater than both. Increase in BS mean CON velocity (m·s ⁻¹) 80% 1RM: TEMPO 1 vs TEMPO 2 (9.85%) and TEMPO 3 (17.65%), TEMPO 1 significantly greater than both. Increase in BS peak CON velocity (m·s ⁻¹) 80% of 1RM: TEMPO 1 vs TEMPO 2 (9.71%) and TEMPO 3 (10.04%), TEMPO 1 significantly greater than both. Increase in BP mean CON velocity (m·s ⁻¹) 60% 1RM: TEMPO 1 vs TEMPO 2 (10.66%) and TEMPO 3 (1.47%), TEMPO 1 and 3 significantly greater than TEMPO 2. Increase in BP peak CON velocity (m·s ⁻¹) 60% 1RM: TEMPO 1 vs TEMPO 2 (10.03%) and TEMPO 3 (-1.17%), TEMPO 1 and 3 significantly greater than TEMPO 2. Increase in BP peak CON velocity (m·s ⁻¹) 80% 1RM: TEMPO 1 vs TEMPO 2 (10.06%) and TEMPO 3 (3.97%), TEMPO 1 was significantly greater than TEMPO 2.
Pryo [90]	N = 24 males, age 20.7 ± 1.7 yrs, height 176.3 ± 6.7 cm, BM = 84.0 ± 14.1 kg, and BP 1RM 101.5 ± 19.9 kg.	To determine the effects of various eccentric tempo durations on repetition completed and peak and mean power.	Exercise: BP, volume: repetitions till failure, intensity 80% of 1RM. ECC TEMPOS: TEMPO 1: 1/0/X/3, TEMPO 2: 4/0/X/3, TEMPO 3: 1/0/X/0, TEMPO 4: 4/0/X/0 CON phase velocity was maximized, averaging 1.55 ± 0.3 s across all tempo trials.	Greater increase in repetition completed for TEMPO 1 vs TEMPO 2 (70%) and TEMPO 3 vs TEMPO 4 (69.09%), both TEMPO 1 and 3 significantly greater. Increase in peak power (W) for TEMPO 1 vs TEMPO 2 (26.47%) and TEMPO 3 vs TEMPO 4 (19.44%) both TEMPO 1 and 3 were significantly greater. Increase in mean power (W) for TEMPO 1 vs TEMPO 2 (25.15%) and for TEMPO 3 vs TEMPO 4 (19.49) both TEMPO 1 and 3 were significantly greater.
Wilk [71]	N = 20 healthy female athletes, age = 27.3 ± 2.2 yrs; BM = 53.3 ± 7.7 kg, TE= 3.9 ± 0.63 yrs, close grip BP 1RM = 55.2 ± 9.5 kg, wide grip BP 1RM = 52.7 ± 8.5 kg.	To determine the effects of different ECC tempos on close and wide grip BP repetitions to failure.	Exercises: close grip BP and wide grip BP, volume: 5 x CON failure, intensity: 70% of 1RM. ECC TEMPOS: ECC-2s: 2/0/X/0, ECC-6s: 6/0/X/0,	Increase in total reps ECC-2s vs ECC-6s for close (20.99%) and wide grip BP (19.85%), ECC-2s was significantly greater for both. Increase in TUT for ECC-6s vs ECC-2s for close (36.19%) and wide grip BP (28.67%), ECC-6s was significantly greater for both.
Sampson [91]	N = 12 recreational trained males age = 26.0 ± 3.9 yrs, height = 180.5 ± 7.9 cm, and BM = 79.1 ± 11.9 kg.	To determine the effects of different bicep curl techniques on repetition velocity,	Bicep curl via specialised flywheel device on dominant arm. Volume: 1 x repetition to failure with each protocol. ECC-X: X/0/X/0	No difference between groups for total workload (j). Greater amount of TUT (s) between ECC-2s vs ECC-X (18.75%). Significantly greater repetition mean velocity (m·s ⁻¹) for ECC-X vs ECC-2s during the first 3 repetitions

		TUT and total workload.	ECC-2s: 2/0/X/0	
Wilk [92]	N = 90 healthy males age = 25.8 ± 5.3 yrs, BM = 80.2 ± 14.9 kg, TE = 3.9 ± 4.2 yrs.	To determine the effects of different ECC tempos on 1RM BP performance.	BP 1RM ECC-V: V/0/V/0 (voluntary tempo), ECC-2s: 2/0/V/0, ECC-5s: 5/0/V/0, ECC-8s: 8/0/V/0, ECC-10s: 10/0/V/0.	Increase in 1RM score ECC-V vs ECC-2s (0.41%), ECC-5s (6.68%), ECC-8s (13.35%), ECC-10s (14.64%). ECC-V was significantly greater than ECC-5s, ECC-8s and ECC-10s. Increase in 1RM score ECC-2s vs ECC-5s (6.24%), ECC-8s (12.89%), ECC-10s (14.16%), ECC-2s was significantly greater than ECC-5s, ECC-8s and ECC-10s. Increase in 1RM score ECC-5s vs ECC-8s (6.26%) and ECC-10s (7.46%), ECC-5s was significantly greater than ECC-8s and ECC-10s.
Wilk [93]	N = 21 strength trained females, age = 23.4 ± 2.2 yrs, BM = 52.3 ± 6.7 kg, TE: 2.3 ± 1.5 yrs.	To determine the effects of different ECC tempos on 1RM performance.	1RM BP ECC-2s: 2/0/X, ECC-4s: 4/0/X, ECC-6s: 6/0/X.	Increase in BP 1RM score for ECC-2s vs ECC-4s (6.70%) and ECC-6s (9.85%), ECC-2s significantly greater than both. Increase in 1RM score for ECC-4s vs ECC-6s (2.97%) with a greater increase.
Calixto [67]	ECC-0.5s group: n = 8 males, age = 23.6 ± 0.4 yrs, BM = 81.0 ± 4.1 kg, height = 176.2 ± 1.2 cm, TE = 4.8 ± 0.9 yrs, 1RM ECC BS = 132.2 ± 7.7 kg. ECC-3s group: n = 8 males, age = 26.5 ± 0.6 yrs, BM = 76.3 ± 2.6 kg, height = 175.3 ± 0.8 cm, TE 5.6 ± 1.2 yrs, 1RM ECC BS = 120.8 ± 6.1 kg.	To determine the effects of ECC BS tempo durations on BL GH and total volume of load lifted.	Exercise: BS, volume 4 x 8 ECC only BS for both groups, intensity: 70% of 1RM, ECC TEMPOS: ECC-0.5s: 0.5 s ECC phase, ECC-3s: 3 s ECC phase.	ECC-0.5s had greater total volume of load lifted (9.2%), no difference between groups. Significant increase in BL (mM) during ECC-0.5s from pre to 0, 3, and 6-mins post exercise. Significant increase in BL (mM) during ECC-3s from PRE to 0, 3, 6-, 9-, 15-, and 20-mins post exercise. ECC-3s had greater increase in BL (mM) than ECC-0.5s at 3, 6-, 9-, 15-, and 20-mins post exercise. Peak BL (mM): ECC-3s vs ECC-0.5s (49.18%), ECC-3s was significantly greater than ECC-0.5s. GH (ng·mL ⁻¹ L) post 15 mins: ECC-0.5s (0.1 ± 0.0), ECC-3s (1.7 ± 0.6), ECC-3s was significantly greater than ECC-0.5s.
Wilk [65]	N = 16 male, age = 21-29 yrs BM = 85.9 ± 7.7 kg, BP 1RM = 130.0 ± 17.5 kg, TE = 5.7 ± 1.3 yrs.	The effects of 2 s and 6 s ECC cadence BP on TUT, BL, CK, T and C.	Exercise: BP, volume: 5 x failure, intensity: 70% of 1RM, ECC tempos: ECC-2s: 2/0/2/0 and ECC-6s: 6/0/2/0.	ECC-2s achieved significantly greater volume than ECC-6s. ECC-6s achieved significantly greater TUT, LA, CK, T and C than ECC-2s.
Martins-Costa [94]	N = 15 male age = 24.4 ± 4.4 yrs, height = 177.0 ± 7.0 cm, BM = 78.3 ± 9.4 kg, BP 1RM = 93.4 ± 10.4 kg, TE = >6.0 mon.	To determine the effects of different ECC TEMPOS on IEMG and BL scores during the BP exercise.	Exercise: BP, volume: 3 x 6, intensity: 60% of 1RM, 3 min rest between sets. ECC TEMPOS: ECC-2s: 2 s ECC, 2 s CON. ECC-4s: 4 s ECC, 2 s CON. Blood samples taken after 10 mins of sit-down rest prior to study and 1 minute post set during testing.	ECC-4s vs ECC-2s had significantly greater IEMG Pectoralis major during all 3 sets. ECC-2s and ECC-4s had significantly greater activity during set 2 and 3 compared to set 1. ECC-4s had significantly greater IEMG tricep brachii during all 3 sets. ECC-2s and ECC-4s had significantly greater activity during set 3 compared to set 1.

				<p>ECC-2s and ECC-4s had significantly greater BL (mM) scores from pre to all 3 sets. ECC-4s had significantly greater scores compared to ECC-2s during all 3 sets.</p> <p>ECC-2s and ECC-4s had significantly greater BL (mM) post set 3 compared to 1 and 2.</p> <p>ECC-2s and ECC-6s had significantly greater BL (Mm) post set 2 compared to 1.</p>
--	--	--	--	---

N: Sample size, **YRS:** years, **MON:** Months, **BM:** Body mass, **kg:** Kilograms, **TE:** Training experience, **1RM:** One repetition maximum, **RM:** Repetition maximum, **ECC:** Eccentric, **CON:** Concentric, **TEMPO:** Eccentric/isometric/concentric/isometric, **TUT:** Time under tension, **X:** Fast as possible, **BL:** Blood lactate, **CK:** Creatine kinase, **S:** Seconds, **MINS:** Minutes, **GH:** Growth Hormone, **CK:** Creatine Kinase, **T:** Testosterone, **C:** Cortisol, **BS:** Back squat, **BP:** Bench Press, **iEMG:** integrated electromyography.

Table 4: Acute isokinetic dynamometer studies.

Author	Population	Aim	Method	Findings
Shepstone [53] study part 2.	N = 9 recreationally active (no weight training and no more than 1 structured exercise bout per week) males age = 23.2 ± 2.4 yrs, height = 181.9 ± 6.1 cm, BM = 81.1 ± 5.6 kg.	To determine the effects of 210°·s ⁻¹ and 20°·s ⁻¹ ECC IKD on Z line streaming.	Exercise: ECC IKD maximal arm lengthening. Volume: 3 x 10, rest: 3 min rest between sets. IKD testing values: 210°·s ⁻¹ and 20°·s ⁻¹ . Z-line streaming obtained from muscle biopsy.	ECC-210°·s ⁻¹ and ECC-20°·s ⁻¹ significantly increased z-line streaming, ECC-210°·s ⁻¹ was significantly greater than ECC-20°·s ⁻¹ . No significant increase in Extreme – moderate for both groups. Z-band streaming. (encompassing >10 serially or longitudinally adjacent sarcomeres with damage).
Roschel [54]	ECC-210°·s ⁻¹ group n = 9 males, age = 26.4 ± 4.3 yrs, height = 177.0 ± 3.0 cm, BM = 76.3 ± 9.6 kg. ECC-20°·s ⁻¹ group n = 11 males, age = 25.4 ± 5.0 yrs, height = 176.0 ± 6.0 cm, BM = 77.2 ± 10.5 kg.	To determine the effects of 210°·s ⁻¹ and 20°·s ⁻¹ ECC IKD on total work, peak torque, total protein, and protein phosphorylation, mTOR, p70 ^{S6k1} , Mechano growth factor, mRNA, Akt phosphorylation expression, mTOR phosphorylation and p70 ^{S6k1} .	Exercise: ECC IKD Knee extension (0° to 90°). Volume: 5 x 8, rest: 3 min rest between sets. Testing values: 210°·s ⁻¹ and 20°·s ⁻¹	No difference between groups for: Total work, peak torque total protein and protein phosphorylation, Mtor, p70 ^{S6k1} . Significant increase for ECC-20°·s ⁻¹ group in baseline and 2 hours post exercise for; Mechano growth factor and mRNA expression. Significant increase for both groups in; Akt phosphorylation from baseline to immediately post and 2 hours post exercise, mTOR phosphorylation immediately after exercise and p70 ^{S6k1} immediately after exercise and post 2 hours exercise.
Cress [25]	N = 30 females: age = 25.0 ± 2.6 yrs.	To determine the effects of various IKD torque values to determine changes in muscle force during leg extension.	Exercise: ECC IKD (right) leg extension, testing angular velocities: 30, 60, 90, 120, 150, 180, and 210°·s ⁻¹ .	No significant difference between testing values.
Drury [95]	N = 11 healthy males age = 19.8 ± 1.6 yrs, height = 179.7 ± 3.8 cm, BM = 84.2 ± 14.7 kg, TE = >6.0 mon.	To determine the effects of different angular velocities on peak torque during an IKD bicep curl.	Exercise: Bicep curl, extension 170° and flexion 45° Testing torque values: 90, 180, and 300°·s ⁻¹ .	No significant difference between testing torque values.
Hortobágyi [96]	40 males, Group 1: n = 20, age = 22.6 ± 5.0 yrs, BM = 76.6 ± 11.5 kg, height = 178.0 ± 9.0 cm. Group 2: n = 20, age = 23.1 ± 1.8 yrs, BM = 90.9 ± 3.3 kg, height = 178.0 ± 7.0 cm.	To determine the effects of peak torque during IKD ECC arm flexion.	Exercise: Arm flexion and extension Testing values: 30, 90, and 120°·s ⁻¹ . Group 1: low strength group, Group 2: high strength group. Groups were selected by the participant's strength levels.	No significant difference in torque values for group 1. Significant increase for group 2 during 30–120°·s ⁻¹ values for both flexion (8.70%) and extension (6.49%).

--	--	--	--	--

N: Sample size, **YRS:** years, **MON:** Months, **BM:** Body mass, **kg:** Kilograms, **TE:** Training experience, **CM:** Centimetres, **ECC:** Eccentric, $^{\circ}\cdot s^{-1}$: Degrees per second, $^{\circ}$: Degrees, **MINS:** Minutes, **IKD:** Isokinetic dynamometry.

Table 5: Chronic tempo training studies.

Author	Population	Aim	Method	Findings
Mike [97]	N = 30 males ECC-2s: age = 22 ± 2.1 yrs, height = 180.0 ± 6.6 cm, BM = 79.0 ± 5.4 kg, TE: 3.1 ± 0.9 yrs, BS 1RM = 124.0 ± 20.0 kg. ECC-4s: age = 22.0 ± 2.1 yrs, height = 176.0 ± 4.8 cm, BM = 82.0 ± 12.0 kg, TE = 2.8 ± 1.1 yrs, BS 1RM = 129.0 ± 22.0 kg. ECC-6s: age = 23.0 ± 4.2 yrs, height = 178.0 ± 8.4 cm, BM = 85.0 ± 16.7 kg, TE = 2.8 ± 1.1 yrs, BS 1RM = 129.0 ± 22.0 kg.	To determine the effects of different ECC TEMPOS on various performance metrics.	Duration: 4-weeks 2 x days per week for 8 sessions. Sessions 1–4, volume: 4 x 6, intensity 80% of 1RM, exercise: BS. Sessions 5–8, volume: 4 x, intensity: 85% of 1RM. Rest: 3 min rest between sets. ECC tempos: ECC-2s: 2/1/2, ECC-4s: 4/1/2, ECC-6s: 6/1/2.	Increase in 1RM BS: ECC-2s (8.87%), ECC-4s (13.18%), ECC-6s (12.71%) no difference between groups. Increase in vertical jump height (inches): ECC-2s (4.5%), ECC-4s (1.4%) and ECC-6s (1.6%), ECC-2s significant increase. Significant decrease for ECC-2s for pre to post study perceived soreness 24, 48 and 72 hrs post training session. Significant decrease for ECC-4s for pre to post study perceived soreness 24 and 48 hrs post training session. Significant decrease for ECC-6s for pre to post study in perceived soreness 48 hrs post training session.
Pereira [69]	Gender of participants not identified. ECC-1s: age = 28.3 ± 8.2 yrs, height = 172.3 ± 5.3 cm, BM = 72.3 ± 9.3 kg, BF% = 17.3 ± 2.2%. ECC-4s: age = 30.3 ± 5.6 yrs, height = 172.6 ± 4.8 cm, BM = 73.8 ± 5.1 kg, BF% = 19.3 ± 0.9%.	To determine the effects of different ECC tempos Scott curls on bicep 1RM and CSA.	Duration: 12 weeks, 2 sessions per week. Volume: 3 x 8 maximum, load not identified. Exercise: Scott curl. ECC TEMPOS: ECC-1s: 1/0/1/0, ECC-4s: 4/0/1/2.	Increase in CSA (cm ²) ECC-1s (6.25%) and ECC-4s (16.55%). Greater increase with ECC-4s vs ECC-1s. Increase in 1RM (kg) ECC-1s (18.98%) and ECC-4s (32.40%). Greater increase with ECC-4s vs ECC-1s.
Stasinaki [68]	N = 18: ECC-1s: n = 5 males 4 females, age = 21.9 ± 0.7 yrs, height 173.4 ± 9.8 cm, BM = 65.2 ± 10.1 kg, TE = 0.0 yrs. ECC-4s: n = 5 males 4 females = age = 22.1 ± 2.0 yrs, height 170.4 ± 7.8 cm, BM = 67.7 ± 9.9 kg, TE = 0.0 yrs.	To determine the effects of different ECC tempos on performance variables.	Duration: 6-weeks, 2 sessions per week. Exercise: ECC only half BS. ECC-1s: ECC TEMPO: <1 s, volume: 9 x 9, intensity: 70% of 1RM, 3 min rest between sets. ECC-4s: ECC TEMPO: 4 s, volume: 5 x 6, intensity 90% of 1RM, 3 min rest between sets.	Increase in BS 1RM Squat ECC-1s (14.5 ± 7.0%) and ECC-4s (5.4 ± 5.1%), ECC-1s was significantly greater than ECC-4s. Increase in CMJ height ECC-1s (2.1 ± 7.6%), ECC-4s (1.3 ± 7.5%) no difference between groups. Increase in CMJ power ECC-1s (7.9 ± 22.2%), ECC-4s (-6.8 ± 7.8%), ECC-4s significantly decreased, no difference between groups. Increase in vastus lateralis muscle thickness: ECC-1s (3.1 ± 9.1%), ECC-4s (6.8 ± 7.8%), ECC-4s was significantly greater, no difference between groups. Increase in fascicle angle: ECC-1s (5.7 ± 15.5%), ECC-4s (7.8 ± 18.8%), no difference between groups. Increase in fascicle length: ECC-1s (10.0 ± 6.2%), ECC-4s (-3.1 ± 7.1%), significant increase from pre to post for ECC-1s and ECC-1s vs ECC-4s.
Douglas [43]	N = 14 male rugby academy players age = 19.4 ± 0.8 yrs, height = 182.0 ± 5.0	The effects of 2, 4-week AEL vs TRAD	2 Part training study.	Increase in relative back squat 1RM kg.BM ⁻¹ :

	cm, BM = 97.0 ± 11.6 kg, relative BS 1RM 1.7 ± 0.2 kg.BM ⁻¹ , TE = > 1.0 yrs.	programmes performed at 1 s and 3 s ECC TEMPOS on performance variables.	Week 1–4: ECC TEMPOS: AEL3s: 3 s AEL, TRAD3s: 3 s TRAD. 2-weeks rest between programmes. Week 4–8: AEL1s: 1 s AEL, TRAD1s: 1 s TRAD eccentric phase.	AEL-1s (-0.54%), AEL-3s (5.03%), TRAD-1s (0.63%), TRAD-3s (-1.82%). 0.50-m drop jump variables Increase in contact time (s): AEL-1s (-9.09%), AEL-3s (0.00%), TRAD-1s (0.00%), TRAD-3s (0.00%). Increase in flight time (s): AEL-1s (-2.04%), AEL-3s (0.00%), TRAD-1s (0.00%), TRAD-3s (2.22%). Increase in RSI: AEL-1s (5.54%), AEL-3s (0.00), TRAD-1s (0.51%), TRAD-3s (1.03%). Leg stiffness (kM.m.kg ⁻¹) AEL-1s (17.86%), AEL-3s (0.00%), TRAD-1s (4.00) TRAD-3s (-3.85%). Increase in muscle thickness (cm): AEL-1s (3.23%), AEL-3s (3.33%), TRAD-1s (3.12%), TRAD-3s (0.00%). Increase in fascicle angle (θ): AEL-1s (5.04%), AEL-3s (-3.47%), TRAD-1s (3.31%), TRAD-3s (0.00%). Increase in Fascicle length (cm): AEL-1s (-6.77%), AEL-3s (9.02%), TRAD-1s (-0.80%), TRAD-3s (-0.79%).
Shibata [98]	ECC-2s group: n = 11, age = 19.8 ± 0.9 yrs, height = 170.7 ± 3.1 cm, and BM = 65.6 ± 4.9 kg. ECC-4s group: n = 11, age = 19.9 ± 1.0 yrs, height = 173.6 ± 4.0 cm, BM = 66.6 ± 8.1 kg.	To determine the effects of different ECC TEMPOS on performance variables.	Duration: 6 weeks 2 x sessions per week. Exercise: BS, volume: 3 x failure, intensity: 75% of 1RM, ECC TEMPOS: ECC-2s: 2 s ECC, 2 s CON ECC-4s: 4 s ECC, 2 s CON	No difference in TUT and volume between 2 s and 4 s. 2 s and 4 s increase in CSA at distal, proximal, and mid, no difference between groups. Increase in BS 1RM ECC-2s (18.46%) and ECC-4s (10.46%), greater increase in ECC-2s vs ECC-4s. Increase in CMJ and SJ jump height, no difference between ECC-2s vs ECC-4s. No change in T-Test and YoYo run for both groups.
Kojić [70]	N = 11 males 9 females. ECC-1s group: age = 24.5 ± 2.2 yrs, height = 178.0 ± 7.0 cm, BM = 72.2 ± 13.5 kg. ECC-4s group: age = 23.6 ± 0.9 yrs, height = 172 ± 8.0 cm, BM = 68.6 ± 11.6 kg.	To determine the difference of different ECC tempo on biceps brachii thickness and 1RM.	Duration: 7 weeks, 2 x sessions per week. Exercise: bicep curl. Weeks 1 – 3, volume: 3 x failure, intensity: 60% of 1RM, Weeks 4 – 7, volume: 4 x failure, intensity: 70% of 1RM. ECC TEMPOS: ECC-1s: 1/0/1/0, and ECC-4s: 4/0/1/0.	Increase in bicep brachii thickness: ECC-1s (15.4%), ECC-4s (18.3%), significantly greater increase in ECC-4s vs ECC-1s. Increase in 1RM: ECC-1s (11.6%), ECC-4s (23.5%), significantly greater increase in ECC-4s vs ECC-1s.

N: Sample size, **YRS:** years, **BM:** Body mass, **kg:** Kilograms, **CM:** Centimetres, **TE:** Training experience, **BF%:** Body fat percentage, **1RM:** One repetition maximum, **ECC:** Eccentric, **CON:** Concentric, **TUT:** Time under tension, **TEMPO:** Eccentric/isometric/concentric/isometric, **CSA:** Cross sectional area, **MINS:** Minutes, **S:** Seconds, **BS:** Back squat, **BP:** Bench Press, **CM:** Countermovement jump, **AEL:** Accentuated eccentric loading, **TRAD:** Traditional.

Table 6: Chronic isokinetic dynamometer studies.

Author	Population	Aim	Method	Results
Farthing [56]	ECC-180°·s ⁻¹ : n = 4 males, 9 females, age = 21.9 ± 1.5 yrs, BM = 65.6 ± 3.8 kg, height 167.0 ± 2.0 cm, TE = 1.2 ± 0.7 mon. ECC-30°·s ⁻¹ : n = 7 males 6 females, age = 19.4 ± 0.6 yrs, BM = 74.3 ± 3.0 kg height = 175.0 ± 2.1 cm, TE = 2.3 ± 0.8 mon. CT: n = 2 males, 8 females, age = 22.7 ± 0.9 yrs, 65.2 ± 3.2 kg, height = 169.0 ± 2.6 cm. 2.3 ± 0.8, TE 2.3 ± 1.1 mon.	To determine the effects of ECC IKD at 180°·s ⁻¹ and 30°·s ⁻¹ on performance variables.	Duration: 8 weeks, 3 sessions per week, a total of 24 training sessions. Volume: progressive increase from 2 to 6 sets of 8 maximal repetitions per training session over the first 13 training sessions, 6 sets for training sessions 13-22, and a taper down to 3 sets of 8 maximal repetitions prior to post-testing during training sessions 23 and 24. Training torque values: ECC-180°·s ⁻¹ : 180°·s ⁻¹ and ECC-30°·s ⁻¹ : 30°·s ⁻¹ . 100° range of motion, approximately 60° to 160° elbow extension for eccentric contractions.	Trained arm peak torque (Nm) % increase at ECC 180°·s ⁻¹ : ECC-180°·s ⁻¹ (23.73%), ECC-30°·s ⁻¹ (-5.56%), CT (8.62%). ECC-180 vs CT was significantly greater post study. Trained arm peak torque (Nm) % increase at ECC 30°·s ⁻¹ : ECC-180°·s ⁻¹ (23.21%), ECC-30°·s ⁻¹ (-7.14), CT (8.93%). ECC-180°·s ⁻¹ vs CT was significantly greater post study. Trained arm peak torque (Nm) % increase at CON 180°·s ⁻¹ : ECC-180°·s ⁻¹ (29.73%), ECC-30°·s ⁻¹ (-2.22%), CT (22.58%). ECC-180°·s ⁻¹ and ECC-30°·s ⁻¹ vs CT was significantly greater post study. Trained arm peak torque (Nm) % increase at CON 30°·s ⁻¹ : ECC-180°·s ⁻¹ (34.29%), ECC-30°·s ⁻¹ (-4.26%), CT (11.76%). ECC-180°·s ⁻¹ and ECC-30°·s ⁻¹ vs CT was significantly greater post study+. No significant increase at the proximal site for the trained arm for all groups. Increase in muscle thickness mid-site: ECC-180°·s ⁻¹ (12.50%), ECC-30°·s ⁻¹ (8.56%), CT (-3.13%), Both ECC-180°·s ⁻¹ and ECC-30°·s ⁻¹ significantly increased from pre to post. Increase in muscle thickness distal site: ECC-180°·s ⁻¹ (14.71%), ECC-30°·s ⁻¹ (10.81%), CT (0%). Both ECC-180°·s ⁻¹ and ECC-30°·s ⁻¹ significantly increased from pre to post.
Farthing [55]	ECC-180°·s ⁻¹ group n = 4 males, 9 females, age = 21.9 ± 1.5 yrs, BM = 65.6 ± 3.8 kg, height = 167.1 ± 2.2 cm, TE = 1.2 ± 0.6 MON. ECC-30°·s ⁻¹ group n = 7 males, 4 females age = 19.6 ± 0.7 yrs, BM = 74.6 ± 3.5 kg, height = 175.7 ± 2.3 cm, TE = 2.7 ± 1.1 MON, CT: n = 2 males, 8 females age = 22.7 ± 0.9 yrs height = 168.8 ± 2.6 cm BM 65.2 ± 3.2 kg TE = 2.3 ± 1.1 MON.	To determine the effects of 180°·s ⁻¹ and 30°·s ⁻¹ ECC IKD on performance variables.	Duration: 8 weeks, 3 sessions per week, 24 training sessions. Exercise: elbow flexion, training torque values: ECC-180°·s ⁻¹ : 180°·s ⁻¹ , ECC-30°·s ⁻¹ : 30°·s ⁻¹ and CT. Volume: 2-6 sets x 8 maximal repetitions per training session over the first 13 training sessions. 6 sets for training sessions 13–22 and a taper down to 3 x 8 maximal repetitions prior to post testing (training sessions 23 and 24).	Increase in combined sites (proximal, mid, and distal) muscle thickness: ECC-180°·s ⁻¹ (13%), ECC-30°·s ⁻¹ , (7.8%), CT (-0.8%). ECC-180°·s ⁻¹ and ECC-30°·s ⁻¹ vs CT was significantly greater Significantly greater increase in ECC torque at 180°·s ⁻¹ and 30°·s ⁻¹ for ECC-180°·s ⁻¹ vs ECC-30°·s ⁻¹ and CT. Significantly greater increase in CON torque at 180°·s ⁻¹ and 30°·s ⁻¹ , for ECC-180°·s ⁻¹ vs CT.
Shepstone [53] study 1	N = 12 male age 23.8 ± 3.4 yrs, height = 178.5 ± 9.6 cm, BM = 82.5	To determine the effects of 210°·s ⁻¹ and 20°·s ⁻¹	Duration: 8 weeks, 3 sessions per week. Volume: week 1: 1 x 10, week	Increase in peak torque (N.m) at torque testing values of 20, 60, 120, 180 and 210°·s ⁻¹ during both ECC and CON muscle actions:

	<p>± 10.3 kg. Not engaged in RT 6 MON prior to the study.</p>	<p>ECC IKD on performance variable in a training study.</p>	<p>2: 2 x 10, week 3: 3 x 10, weeks 4–8: 4 x 10 with 180 s rest between sets. Exercise: elbow arm flexion. ECC training torque values: ECC-210°·s⁻¹: 210°·s⁻¹ and ECC-20°·s⁻¹: 20°·s⁻¹</p>	<p>ECC-20°·s⁻¹ and ECC-210°·s⁻¹ increased from pre to post, ECC-210°·s⁻¹ had a greater improvement than ECC-20°·s⁻¹. Increase in CSA from pre to post: ECC-210°·s⁻¹ had significantly greater increase vs ECC-20°·s⁻¹. Increase in muscle fibre size type Ia for pre to post study: both ECC-210°·s⁻¹ and ECC-20°·s⁻¹ significantly increased. Increase in muscle fibre size type IIa for pre to post study: both ECC-210°·s⁻¹ and ECC-20°·s⁻¹ significantly increased, ECC-210°·s⁻¹ vs ECC-20°·s⁻¹ had a greater increase. Increase in muscle fibre size type IIx for pre to post study: both ECC-210°·s⁻¹ and ECC-20°·s⁻¹, significantly increased, ECC-210°·s⁻¹ vs ECC-20°·s⁻¹ had a greater increase. Increase in muscle fibre % pre to post study: Type Ia: ECC-210°·s⁻¹ (-9.81%), ECC-20°·s⁻¹ (10.18%). Type IIa: ECC-210°·s⁻¹ (10.65%) ECC-20°·s⁻¹ (-1.61%). Type IIx: ECC-210°·s⁻¹ (-25.27%) ECC-20°·s⁻¹ (-34.51%), significant decrease for both ECC-210°·s⁻¹ and ECC-20°·s⁻¹. Increase in muscle fibre area % from pre to post study: Type Ia: ECC-210°·s⁻¹ (-16.52%), ECC-20°·s⁻¹ (4.92%). Type IIa: ECC-210°·s⁻¹ (14.43%), ECC-20°·s⁻¹ (2.93%), ECC-210°·s⁻¹ significantly improved from pre to post and significantly greater vs ECC-20°·s⁻¹. Type IIx ECC-210°·s⁻¹ (-34.12%) ECC-20°·s⁻¹ (-34.41%). Increase in myosin heavy chains %: Type Ia: ECC-210°·s⁻¹ (18.77%) ECC-20°·s⁻¹ (-7.84). Type IIa: ECC-210°·s⁻¹ (21.26%) ECC-20°·s⁻¹ (14.79%), both ECC-210°·s⁻¹ and ECC-20°·s⁻¹ significantly increased from pre to post study and ECC-210°·s⁻¹ vs ECC-20°·s⁻¹ was significantly greater. Type IIx: ECC-210°·s⁻¹ (-42.50) ECC-20°·s⁻¹ (-31.34%), both significantly decreased.</p>
<p>Paddon-Jones [58]</p>	<p>N = 20 non-resistance trained males and females age = 24.2 ± 7.0 yrs, height = 176.6 ± 5.9 cm, BM 76.2 ± 15.8 kg. ECC-180°·s⁻¹: n = 5 male 2 female. ECC-30°·s⁻¹: n = 5 male 1 female. CT: n = 5 males 2 females.</p>	<p>To determine the effects on torque and muscle fibre change when completing either 180°·s⁻¹ or 30°·s⁻¹ IKD ECC training.</p>	<p>Duration: 3 non consecutive days for 10 weeks Volume: 4 x 6 repetitions 60 s rest between sets. Exercise: non dominant arm flexion. ECC IKD values: ECC-180°·s⁻¹: 180°·s⁻¹ and ECC-30°·s⁻¹: 30°·s⁻¹</p>	<p>Significant increase in isometric torque for ECC-180°·s⁻¹ group from pre to post and compared to CT group. Increase in CON torque at 30°·s⁻¹: no significant increase for any group from pre to post. Increase in CON torque at 180°·s⁻¹: Significant increase from pre to 5 weeks for both ECC-180°·s⁻¹ and ECC-30°·s⁻¹ groups. Post 10 weeks ECC-180°·s⁻¹ had significantly greater improvement vs ECC-30°·s⁻¹ and CT.</p>

				<p>Increase in ECC torque $30^{\circ}\cdot s^{-1}$: ECC-$180^{\circ}\cdot s^{-1}$ had significantly greater improvement post study vs ECC-$30^{\circ}\cdot s^{-1}$ and CT.</p> <p>Increase ECC torque $180^{\circ}\cdot s^{-1}$: ECC-180 had significantly greater improvement post study vs CT.</p> <p>Increase in muscle fibre changes: ECC-$180^{\circ}\cdot s^{-1}$ significant increase type IIb but significantly decreased type Ia fibre from pre to post. No significant difference between the other groups.</p>
Ünlü [60]	<p>20 males, ECC-$180^{\circ}\cdot s^{-1}$: n = 6, age = 20.5 ± 1.9 yrs, BM = 72.8 ± 8.2 kg, height = 187.8 ± 8.5 cm.</p> <p>ECC-$30^{\circ}\cdot s^{-1}$: n = 7, age = 21.6 ± 0.5 yrs, BM = 68.7 ± 8.2 kg, height = 176.7 ± 4.9 cm. CT: n = 7, 21.3 ± 2.4 yrs, BM = 67.6 ± 8.8 kg, height = 178.4 ± 6.3 cm.</p>	<p>To determine the effects of completing either $180^{\circ}\cdot s^{-1}$ or $30^{\circ}\cdot s^{-1}$ isotonic leg extension during a 12-week programme on performance variables.</p>	<p>Duration: 12-week training programme in respected group. 3 x per week. Exercise: isotonic knee extension.</p> <p>ECC-$180^{\circ}\cdot s^{-1}$: ECC only at $180^{\circ}\cdot s^{-1}$, 100-155% or 1RM and time under load per set 8 – 10 s.</p> <p>ECC-$30^{\circ}\cdot s^{-1}$: ECC only at $30^{\circ}\cdot s^{-1}$, 80-85% or 1RM, time under load per set 30–40 s</p> <p>CT: Non training control group.</p>	<p>Increase in 1RM: ECC-$180^{\circ}\cdot s^{-1}$ (28.5%), ECC-$30^{\circ}\cdot s^{-1}$ (36.5%), CT (-0.6%), ECC-$180^{\circ}\cdot s^{-1}$ and ECC-$30^{\circ}\cdot s^{-1}$ had greater increase than CT.</p> <p>Increase in peak torque CON $60^{\circ}\cdot s^{-1}$: ECC-$180^{\circ}\cdot s^{-1}$ (13.5%), ECC-$30^{\circ}\cdot s^{-1}$ (24.9%), CT (-0.8%), ECC-$180^{\circ}\cdot s^{-1}$ and ECC-$30^{\circ}\cdot s^{-1}$ had a greater increase than CT.</p> <p>Increase in peak torque CON $180^{\circ}\cdot s^{-1}$: ECC-$180^{\circ}\cdot s^{-1}$ (15.10%), ECC-$30^{\circ}\cdot s^{-1}$ (17.00%), CT (3.00%), ECC-$180^{\circ}\cdot s^{-1}$ and ECC-$30^{\circ}\cdot s^{-1}$ had a greater increase than CT.</p> <p>Increase in muscle volume (cm^3): ECC-$180^{\circ}\cdot s^{-1}$ (4.14%), ECC-$30^{\circ}\cdot s^{-1}$ (10.04%), CT (3.75%). No difference between groups.</p>

N: Sample size, **YRS:** years, **BM:** Body mass, **kg:** Kilograms, **TE:** Training experience, **M:** Meters, **CM:** Centimetres, **ECC:** Eccentric, **CON:** Concentric, $^{\circ}s^{-1}$: Degrees per second, $^{\circ}$: Degrees, **HRS:** Hours, **MINS:** Minutes, **MON:** Months, **CT:** Control, **1RM:** One Repetition Maximum.

Table 7: Isokinetic dynamometry angular velocity values across studies which employ the terms of fast and slow.

Study	Training method	Classified (fast)	Classified (slow)
Shepstone [53] Roschel [54]	IKD	210°·s ⁻¹	20°·s ⁻¹
Farthing [55] Farthing [56] Paddon-Jones [57] Paddon-Jones [58]	IKD	180° s ⁻¹	30° s ⁻¹
Chapman [50]	IKD	210°·s ⁻¹	60°·s ⁻¹
Aagaard [59]	IKD	240°·s ⁻¹	60°·s ⁻¹
Ünlü [60]	Isotonic training	180°·s ⁻¹	30°·s ⁻¹

IKD: Isokinetic dynamometry; °·s⁻¹: Degrees per second; Note: All values were calculated to °·s⁻¹ for comparison, calculated via Excel with the following formula 1 rad·s⁻¹ = 57.3°·s⁻¹.

Table 8: Movement tempo durations for various studies which use the term of fast and slow.

Study	Exercise	Defined as eccentric or overall tempo	Classified (fast)	Classified (slow)
Wilk [65]	Bench Press	Eccentric	Regular: 2 s	Slow: 6 s
Wilk [66]	Bench Press	Eccentric	Regular: 2 s	Slow: 6s
Calixto [67]	Bench Press	Eccentric	Fast: 0.5 s	Slow: 3 s
Douglas [43]	Back Squat	Eccentric	Fast: 1 s	Slow: 3 s
Stasinaki [68]	Back Squat	Eccentric	Fast: X	Slow: 4 s
Pereira [69]	Scott Curl	Eccentric	Fast 1 s	Slow: 4 s
Kojić [70]	Bicep curl	Eccentric	Fast: 1 s	Slow: 4 s
Wilk [71]	Bench Press	Overall	Fast 2/0/X/0,	Slow: 6/0/X/0
Lopes [72]	Bench Press	Overall	Fast: 0.75 s ECC, 0.75 s CON, (total 1.5 s)	SLOW 3 s ECC, 3 s CON, (total 6 s)

REG: Regular, **X:** Fast as possible, **S:** Seconds, **ECC:** Eccentric, **CON:** Concentric, **FAST:** Authors defined this tempo as fast within their research, **SLOW:** Authors defined this as a slow tempo within their research, **REGULAR:** Authors defined this as regular which falls in between the paradigm of fast and slow tempos.




Figure 1: Schematic of search strategy process.